

The Evolving Neural and Genetic Architecture of Vertebrate Olfaction

Daniel M. Bear¹, Jean-Marc Lassance², Hopi E. Hoekstra², and Sandeep Robert Datta^{1,*}

¹Department of Neurobiology, Harvard Medical School, Center for Brain Science, Harvard University, Boston, MA 02115, USA

²Departments of Molecular and Cellular Biology and Organismic and Evolutionary Biology, Center for Brain Science, Harvard University, Howard Hughes Medical Institute, Cambridge, MA 02138, USA

*Correspondence: srdatta@hms.harvard.edu

<http://dx.doi.org/10.1016/j.cub.2016.09.011>

Evolution sculpts the olfactory nervous system in response to the unique sensory challenges facing each species. In vertebrates, dramatic and diverse adaptations to the chemical environment are possible because of the hierarchical structure of the olfactory receptor (OR) gene superfamily: expansion or contraction of OR subfamilies accompanies major changes in habitat and lifestyle; independent selection on OR subfamilies can permit local adaptation or conserved chemical communication; and genetic variation in single OR genes can alter odor percepts and behaviors driven by precise chemical cues. However, this genetic flexibility contrasts with the relatively fixed neural architecture of the vertebrate olfactory system, which requires that new olfactory receptors integrate into segregated and functionally distinct neural pathways. This organization allows evolution to couple critical chemical signals with selectively advantageous responses, but also constrains relationships between olfactory receptors and behavior. The coevolution of the OR repertoire and the olfactory system therefore reveals general principles of how the brain solves specific sensory problems and how it adapts to new ones.

Introduction

Olfaction, the sense of smell, has evolved to detect signals from the chemical environment, which contains clues about where to move, what to eat, when to reproduce and which stimuli to remember as rewarding or dangerous [1,2]. How an animal responds to chemical cues is in part learned and in part innate, depending on how the olfactory nervous system has been shaped by both experience across a lifetime and evolution across generations.

Interacting with the chemical world presents a challenge unlike other sensory tasks. In contrast to light and sound waves, which vary continuously in wavelength and amplitude, chemical compounds differ discretely in an enormous number of dimensions. Compared to vision and audition, olfaction requires many more receptors — proteins expressed in sensory organs that convert a physical event into an electrochemical signal carried by neurons. A single photoreceptor pigment may interact with a range of wavelengths that is sufficiently informative for an animal, as with our dark-vision rhodopsin. However, biophysical constraints restrict chemoreceptor proteins to interact with only subsets of chemical space. The high dimensionality of chemical space and the diversity of the chemical stimuli an animal might encounter have therefore selected for genomes that encode enormous receptor repertoires, containing hundreds to thousands of OR genes [3–6].


Here, we examine how the olfactory receptor repertoire and the structure of the olfactory nervous system evolve in concert to sense and interpret chemical information. Adaptations in the genetic and neural architecture of olfaction reflect the unique chemosensory challenges faced during the evolution of a species — detecting novel odorants, discerning especially critical stimuli with high acuity and responding behaviorally to molecules that acquire new meaning. We argue that the

receptors underlying vertebrate olfaction possess two properties essential for the range of adaptations seen in vertebrate olfactory systems: first, a flexible and hierarchical pattern of evolution that allows receptor adaptation to both dramatic and subtle changes in the chemical environment; and second, expression within a diverse array of neural pathways that govern both hard-wired, instinctual behaviors as well as more flexible odor learning. In this way, OR families and subfamilies have been sculpted to inform the animal about wide swaths of its chemical environment, while at the same time some individual receptors have become highly tuned to key odorants that elicit innate responses. We speculate that the unique ability to incorporate new and evolving OR genes has produced the substantial genetic and structural diversity of olfactory systems seen across vertebrates [7–9].

Molecular, Cellular and Neural Architecture of Vertebrate Olfaction

Smell begins when odor molecules bind to OR proteins on the endings of sensory neurons. The set of odors an animal can detect therefore depends on the expression pattern and the protein structure encoded by each of its OR genes. In vertebrates, nearly all OR genes encode seven-pass transmembrane G-protein coupled receptors (GPCRs) that, upon ligand binding, signal through G-proteins and intracellular second messengers to open membrane ion channels. This depolarizes the sensory neuron to drive action potentials that are conducted along its axon into the olfactory bulb of the brain (Figure 1) [10].

In vertebrates, most olfactory sensory neurons express a single olfactory receptor gene from one of five major GPCR families [11]. The evolutionary history of each family relates roughly to the region of chemical space it probes, as the odorant structures bound by family members are reminiscent of those bound by the specific ancestral, non-olfactory GPCR from which each



Current Biology

Figure 1. Organization of the mouse olfactory system.

Odors are blends of molecular compounds inhaled into the nasal cavity, where they interact with olfactory receptor proteins expressed in the main olfactory epithelium (medium gray), the vomeronasal organ (dark gray), or one of several smaller sensory structures not pictured (top). Each sensory neuron expresses a single olfactory receptor denoted by its color, and neurons expressing the same receptor project to insular structures in the olfactory bulb called glomeruli (bottom). Glomeruli corresponding to a given receptor have stereotyped spatial positions across animals. Mitral cells in the olfactory bulb each send a dendrite into one glomerulus (main olfactory bulb) or multiple glomeruli corresponding to the same olfactory receptor (accessory olfactory bulb, not pictured), and project axons to a variety of central brain regions that mediate odor learning and innate odor-driven behaviors. The targets of main and accessory bulb mitral cell projections are largely distinct (modified after [76]).

family is derived. In the rodent olfactory system, members of each OR family are expressed pointillistically within spatially restricted zones of the nasal epithelia. Most ciliated sensory neurons of the main olfactory epithelium express receptor genes of the largest family, the classical mammalian olfactory receptors (mORs) [12,13], while a minority of neurons instead express members of the much smaller trace amine-associated receptor (TAAR) family [14]. Rodents and most tetrapods also have an ‘accessory’ olfactory system served by the vomeronasal organ, a separate sensory epithelium that detects water-soluble pheromones and other labile chemical signals drawn into the nasal cavity. Microvillar sensory neurons in the apical epithelial layer of the vomeronasal organ express Vomeronasal type 1 Receptors (V1Rs) [15], but neurons in the basal layer each express one Vomeronasal type 2 Receptor (V2R) plus a chaperone V2R in characteristic combinations [16–18]. Finally, some neurons in both layers of the vomeronasal organ instead express members of the small Formyl Peptide Receptor-related family (FPRs) [19,20] (Figures 1 and 2).

The one-receptor-per-neuron organization determines how odor representations propagate through the olfactory system. The axons of primary sensory neurons expressing the same receptor gene coalesce onto insular structures called glomeruli in the outer layer of the olfactory bulb, with the positions of the glomeruli corresponding to a given OR being approximately fixed between individuals (Figure 1) [21]. A given odorant binds

to a stereotyped subset of olfactory receptors and thereby activates a characteristic subset of glomeruli, such that each odor elicits a unique spatiotemporal activity pattern across the olfactory bulb [22].

Odor responses in the olfactory bulb drive innate and learned behaviors via neural projections to more central brain regions. The largest of these bulbar targets is the piriform cortex, a three-layered paleocortical structure in which odor identity is encoded in the patterned activity of distributed neuronal ensembles. Odor representations are less well understood in other targets of the olfactory bulb, including the cortical amygdala and the olfactory tubercle, a subdivision of the ventral striatum. Mitral and tufted cells of the main olfactory bulb each send a single dendrite into one glomerulus and extend parallel axon branches into each of these regions in characteristic patterns, which is thought to underlie the distinct roles of each of these areas in olfactory processing. Spatially diffuse projections to piriform cortex orchestrate the generation of odor-specific neural ensembles that appear optimized for odor learning [23,24]. In contrast, the cortical amygdala receives stereotyped and spatially restricted innervation from individual glomeruli and exhibits spatially biased responses to innately relevant odors; consistent with this apparent hardwiring, the cortical amygdala mediates instinctual responses to some predator odors (Figure 1) [23,25]. In contrast, mitral cells in the accessory olfactory bulb each extend a dendrite into multiple glomeruli (often corresponding to the same vomeronasal receptor) and project, by way of the medial amygdala, to hypothalamic nuclei involved in reproductive and stress responses (Figure 1) [26]. The targets of main and accessory bulb mitral cells are mostly distinct, and thus higher-order olfactory brain regions have likely adapted to receive information about odors sensed by different receptors that trigger distinct behaviors.

Olfactory brain regions, especially the olfactory bulb, vary tremendously in relative size across vertebrate species, and in many vertebrates, the accessory olfactory bulb and the vomeronasal organ are absent [7,8]. Rather than extensively reviewing the comparative anatomy of olfactory systems, we instead focus on the conserved genetic architecture of vertebrate olfaction, which is likely to have arisen at the latest in the common ancestor of all ray-fined fishes and tetrapods (which lived about 460–430 million years ago). The genome of the zebrafish contains multiple members of four of the five major GPCR OR families (ORs, TAARs, V2Rs, and several V1Rs; Figure 2) [27–29] and its sensory neurons generally express these receptors in the same ‘one receptor, one sensory neuron’ pattern as in mice [30–32]. Zebrafish sensory neurons project to insular glomeruli within the olfactory bulb, where innervation from different sensory neuron types (expressing distinct OR families) is at least partially segregated [33,34]. The conservation between zebrafish and mice (as well as many other species) of this convergence from sensory neurons to glomeruli suggests that the neural organization of the olfactory system forms a stable background on which to compare changes in the OR gene repertoire. The molecular mechanism for choosing a single OR to be expressed in each sensory neuron, which engages both cell type-specific transcription factors as well as stochastic DNA–DNA interactions [11], seems to have directly linked the genetic evolution of OR families with the functional evolution of odor




Figure 2. Phylogenetic relationships of the vertebrate GPCR olfactory receptor repertoire.

A representative sample of ~2700 GPCR functional OR genes from zebrafish (*Danio rerio*), frog (*Xenopus laevis*), mouse (*Mus musculus*), and human (*Homo sapiens*) genomes was used to construct a phylogenetic tree [10]. Notable features are visible, including large expansions of Class II ORs (red) in tetrapods, V2Rs (purple) in amphibians, TAARs (blue) in fishes, and V1Rs (teal) in mammals; significant losses of functional ORs in humans (blue dots) compared to mice (green dots); the monophyly of Class I (orange) and Class II (red) tetrapod ORs; and the subfamily structure of the classical ORs, with numeric labels corresponding to the subfamilies defined in [52]. A Class II OR subfamily (2,13) itself contains a smaller subfamily, the “OR37” receptors, which has atypically expanded and been conserved at the protein sequence level in humans. Note that all mammalian and some frog Class I ORs form a monophyletic clade within the larger set of Class I ORs and other fish OR subfamilies that are neither Class I nor Class II.

perception. This process has allowed the olfactory system to flexibly incorporate or discard chemosensors by endowing each functional OR gene with its own population of sensory neurons in the olfactory epithelium and its own glomerular territory in the olfactory bulb. That is, because newly evolved ORs do not directly alter the tuning or processing of existing neural pathways, the functional constraints on receptor gene evolution may be looser for olfaction than for other sensory systems.

Birth-and-Death Mechanism of OR Gene Family Evolution

Singular OR expression potentially affords each newly arising OR gene a neural pathway to inform the brain about the odorants it binds. The evolution of the OR gene repertoire should therefore reflect the salient features of an animal’s chemical environment. To test this hypothesis, genomic observations of OR evolution must be combined with knowledge of what molecules individual receptors bind and how different animals respond to these odors. Unfortunately, because of the enormous number of OR proteins and the difficulty of measuring their molecular tuning *in vitro*, we know only a small fraction of OR–odorant interactions [35].

Nevertheless, due to the mechanism of OR repertoire evolution, it is possible to compare branches of the OR gene tree as they grow or die off on multiple evolutionary timescales. The low proportion of orthologous receptors found among even closely related species suggests that variations in the OR repertoire can emerge and become fixed in relatively short spans of time. This rapid rate of receptor evolution is the result of a characteristic ‘birth-and-death’ process. Changes in the number of OR genes arise from duplications or deletions of contiguous family members, creating redundant and functionally independent new genes that may mutate and acquire new ligand-binding properties [36]. A side-effect of the relaxed selection pressure on recently duplicated genes is the accumulation of nonsense or frameshift mutations that terminate receptor function, producing pseudogenes embedded in rapidly evolving OR gene clusters [37]. This birth-and-death process further explains why few OR genes have true one-to-one orthologous relationships across multiple species, as an ancestral gene may expand into a subfamily of dozens in one lineage in which it finds utility but become pseudogenized or removed in another [38]. Other modes of gene-family evolution have occasionally homogenized the OR repertoire, such as gene conversion between tightly-linked subfamily members and across-species conservation of single ORs that function outside the olfactory system [38–40]. However, it is clear that birth-and-death and relaxed selection are the dominant processes by which different species evolve markedly different sets of olfactory receptors [41,42]. The birth-and-death mechanism has allowed natural selection to independently shape the size and function of different branches of the OR gene tree, including each of the five olfactory GPCR families, more recently diverged subfamilies within each OR family, and individual OR genes (Figure 2).

Widespread Reshaping of the OR Repertoire in New Sensory Environments

As some vertebrates have adapted to new habitats or shifted to lifestyles that rely less on smell, entire OR families and olfactory sensory organs have become dispensable. For example, as their visual system has expanded dramatically, old-world monkeys have seen a massive loss of functional OR genes and a near total loss of the vomeronasal receptor repertoire, accompanied by pseudogenization of an important vomeronasal organ chemotransduction channel gene, *TrpC2* (Figure 2) [43,44]. Selection on primate OR genes may have been relaxed following the duplication of the opsin gene encoding the long-wavelength retinal photopigment, as the genomes of trichomat primates — all old-world monkeys and one species of new-world monkey, the howler monkey — have significantly higher fractions of pseudogenized OR genes than dichromat new world monkeys [45]. Trichromacy may have allowed color vision to take over some olfactory tasks, such as discriminating the ripeness of leaves by color [46,47]. This view is controversial, however, because the proportion of OR pseudogenes in a mammalian genome does not correlate with the number of intact ORs and thus poorly reflects olfactory function [38]. Moreover, loss of intact ORs in primates may have begun before the divergence of new-world monkeys and old-world monkeys [48]. Nevertheless, the growth of the visual system in old-world primates is likely to have reduced reliance on olfaction for many social and foraging

behaviors, which may represent a general pattern: in multiple cases, the enhancement of other senses such as vision, echolocation and electroreception coincides with a smaller OR repertoire [49,50].

Changes in habitat that broadly alter chemical ecology appear to have also resized and reshaped the OR repertoire dramatically, though directly demonstrating causal relationships remains a challenge. The divergence of the tetrapod lineage from its aquatic relatives about 420 million years ago was followed by an order-of-magnitude expansion in the major vertebrate OR gene family (represented by the mORs in mammals). Hundreds to thousands of these classical OR genes are found in individual amphibian and terrestrial vertebrate genomes, whereas fish genomes have at most 150 (Figure 2) [51]. The classical OR gene family has two major branches, the Class I and Class II ORs [52]. However, only the Class II branch shows a massive increase in membership as a possible result of living on land. This receptor subfamily is expanded in the genomes of tetrapods but is essentially absent in aquatic vertebrates, including one more closely related to tetrapods than teleosts — the coelacanth (Figure 2) [53,54].

The idea that having more OR genes is adaptive on land is also strengthened by evidence from several mammal groups that returned to aquatic life, such as whales, seals and manatees. In these animals, the number of intact OR genes has dropped drastically since they diverged from their terrestrial relatives [52].

Why might a particular OR subfamily have expanded in a given vertebrate lineage? The simplest hypothesis is that the ancestral family members were useful for sensing a type of molecular structure that was common or critical in an animal's chemical environment. Having more OR genes with somewhat diversified ligand-binding regions could have provided a selective advantage — for instance by increasing perceptual range, sensitivity or discriminative power in this region of chemical space. The evolutionary history and functional tests of the distinction between Class I and Class II ORs support this model. First identified in the amphibian *Xenopus laevis* genome, Class I genes are closely related to fish ORs, while Class II genes are more closely related to the majority of mammalian ORs (Figure 2) [55]. Furthermore, Class I genes are expressed exclusively in the lateral diverticulum of the frog nasal cavity, which senses waterborne but not airborne odorants from the outside world, while Class II genes are expressed only in the medial diverticulum, which opens only when the frog is in air [55]. Functional characterization supports the hypothesis that *Xenopus* Class I genes are 'fish-like', as they detect highly water-soluble ligands such as charged amino acids, which are important water-borne chemosignals for fish. Conversely, Class II *Xenopus* ORs detect some airborne odors such as the aromas of plants [56]. The ligands of mammalian Class I ORs are also on average more polar (and thus more water-soluble) than Class II ligands [35], but the relative size of the Class I repertoire has remained stable across mammals, suggesting that even in purely terrestrial environments these receptors bind odorants of ecological significance [38].

Particular Habitats May Have Favored Expansion of Distinct OR Families

The presence of smaller "classical" OR families in aquatic animals does not necessarily mean that olfaction is less important

in water than in air. Similar to the above cases of Class I and II ORs, the aquatic environment may have instead favored the use of other OR families because their ancestral ligand-binding properties gave them a head-start in detecting common aqueous odorants. For instance, the TAAR family has expanded rapidly in teleost fishes (Figure 2). The zebrafish genome encodes over 100 functional TAARs, almost an order of magnitude more than any other vertebrate group. This expansion includes an entirely new TAAR subfamily under strong positive selection in the teleost lineage, which was first thought not to encode amine receptors (like the other TAARs) because a canonical amine-binding motif is lost in many subfamily members [27]. It has now been shown, however, that these teleost TAARs initially evolved a second amine-binding motif, changing their specificity to diamines over monoamines. Later, a subset of these secondarily lost the original motif, resulting in TAARs that use a distinct mechanism for amine binding [57]. Together, this pattern of TAAR duplication and diversification suggests that the outsized importance of amine chemosensation drove this particular OR family to prominence in the teleost lineage. This notion is consistent with the important role of amino acids and their diamine decarboxylation products in the chemical ecology of zebrafish [58].

The ancestral TAAR was probably the orthologue of the mammalian TAAR1 and therefore a receptor for internally produced trace monoamines, such as the TAAR1 ligands tyramine, octopamine and tryptamine [59]. Thus, the expansion of the TAAR family in teleosts may have been biased from the start, since only a few mutations in new TAAR genes would allow better sensing of amines, a region of chemical space apparently important for the behavior of these fishes. For example, cadaverine, a diamine arising from putrefaction, is detected by the zebrafish-specific receptor TAAR13c and elicits innate aversion [58]. This link between a novel receptor and amine-driven behavior may be a lineage-specific adaptation, as another teleost, the goldfish, is attracted to cadaverine (perhaps via distinct TAAR receptors; Figure 3) [57–60].

The evolutionary trajectory of the TAARs raises the possibility that, along different lineages, specific ancestral receptors evolved into OR families because they detected molecular structures with wider ecological significance. This reasoning could explain why the V2R family expanded dramatically and reached its largest size in the amphibian lineage (Figure 2) [61]. The V2Rs are derived from an ancestral GPCR that senses the amino acid glutamate and have diversified to allow broader detection of amino acids and longer polypeptides [17]. The amphibian V2R family is expressed in the vomeronasal organ, which plays an essential role in sensing waterborne peptide sex signals and coordinating complex mating behavior [62,63].

These examples suggest that the ecological importance of a given class of chemicals is reflected by the size of the OR family that detects it. It is intriguing to consider the inverse: whether the apparent expansion of a specific OR family can reveal the types of chemical cue that are most relevant to a given species. Consider the case of the platypus, one of only two extant monotremes, which was initially thought not to rely on strong olfactory abilities given the high proportion of pseudogenes in its classical OR repertoire and its use of a unique electroreceptive bill for locating prey in brackish water. However, the platypus genome




Figure 3. Monomolecular odorants that drive innate, species-specific behaviors across vertebrates and insects.

Instinctual reproductive, protective and feeding behaviors are triggered by a variety of chemical structures, including volatile airborne small molecules in rodents (top row); waterborne acids, bases, and peptides in fishes and amphibians (middle row); and volatile hydrocarbons in insects (bottom row). Chemical structures (receptor responsible for behavior, if known), left to right: top row 2,4,5-trimethylthiazoline, 2-sec-butyl-4,5-dihydrothiazole (SBT), 2,3-dehydro-exo-brevicomin (DHB) [76], phenethylamine (*Mus musculus* TAAR4) [84], trimethylamine (*Mus musculus* TAAR5) [85]; middle row 4-hydroxyphenylacetic acid (*Danio rerio* ORA1 (a V1R)) [95], cadaverine (*Danio rerio* TAAR13c) [58], splendiferin (peptide pheromone for *Litoria splendida*), sodefrin (peptide pheromone for *Cynops pyrrhogaster*) [63]; bottom row methylhexanoate (*D. sechellia* Or22a), octanoic acid [78], *cis*-11-vaccenyl acetate (*D. melanogaster* Or67d) [79], (10*E*,12*Z*)-hexadeca-10,12-dien-1-ol (upper) and (10*E*,12*Z*)-hexadeca-10,12-dien-1-ol (lower) (aliases bombykal, receptor *Bombyx mori* BmOr3, and bombykol, receptor *B. mori* BmOr1, respectively) [80]. Behavioral responses are sometimes enhanced by combinations of odorants, as with SBT and DHB in mice and bombykal and bombykol in the silk moth. Moreover, responses to the same compound(s) may differ drastically between species or between sexes of the same species.

contains more functional V1R family members than any other vertebrate, reflecting an unparalleled expansion of this particular gene family in the monotreme lineage [64]. Indeed, the platypus possesses one of the most complex vomeronasal organs described, suggesting that odor detection through the accessory olfactory system has become its main mode of chemosensation. Because V1Rs detect many steroid hormones that regulate reproductive and mating behavior [65], investigating these chemicals may reveal cues critical to the behavior of the platypus and the reasons this particular receptor family has expanded.

Olfactory Specialization through OR Subfamily Selection

OR birth and death operates locally within the genome, producing closely-related receptor subfamilies (Figure 2) [5,37,42] that may evolve independently. There is indirect evidence that natural selection on these finer-scale components of the OR repertoire has allowed each species to better sense the chemical cues relevant to its particular ecological niche. For example, the relative sizes of each mammalian OR subfamily are more representative of a species' ecotype — aquatic, semi-aquatic, terrestrial or flying — than of its ancestry within the phylogenetic tree of mammals [52], suggesting that particular receptor subfamilies have evolved in response to the odors characteristic of each environment.

Testing this hypothesis requires identifying the odors recognized by these receptors and determining whether larger subfamilies actually confer a selective advantage in discerning these cues. Observations of subfamily selection on shorter timescales, correlated with even finer niche adaptation, may help hone in on

the sensory demands driving receptor evolution. For example, in an analysis of OR evolution in bats, the relative sizes of two subfamilies correlate positively and negatively with independent adaptations to a purely frugivorous diet [66]. A similar investigation of bird and reptile OR repertoires identified a subfamily associated with carnivory and several Class I and Class II subfamilies enlarged in aquatic and terrestrial feeders, respectively [67]. Members of these OR subfamilies may have become tuned to volatile compounds found in the foods sought by individual species, and their expression in the olfactory system may reveal how selective neural pathways adapt to influence foraging and food choice.

Similarly, distinct subfamilies of the mouse V2Rs appear tuned to labile, innate fear-producing chemosignals (kairomones) derived from different types of mouse predator [65]. The observation that the family of Major Urinary Proteins (MUP) acts through V2Rs to control both interspecies defensive behaviors and intra-species aggressive behaviors suggests that V2R–MUP pairing or binding affinities may be particular targets of lineage-specific subfamily expansion and selection [68–70]. Furthermore, if different V2R subfamilies are in fact tuned to the chemical signatures of particular predators, their sensory signals could remain segregated as they propagate through the nervous system, which might allow unique defensive responses to evolve for each.

Finally, the modular nature of receptor gene evolution allows small subfamilies to retain critical chemosensory roles, even against a backdrop of widespread receptor loss. For example, the mammalian OR37 subfamily is surprisingly conserved in mice and humans, even though humans have lost more than

half of their functional OR genes overall (Figure 2) [50,71]. Indeed, this unusual subfamily has expanded independently in both the human and the rodent lineage, suggesting it has not lost its utility like other human subfamilies (Figure 2) [72]. Together, these data suggest that humans may still detect a set of informative odors through the OR37 family, such as the long-chain fatty aldehyde ligands sensed by mouse OR37 members [73]. Intriguingly, mouse mitral cell projections from OR37 glomeruli target several hypothalamic nuclei rather than the cortical regions typical of most main olfactory bulb projections [74,75]. Thus, the OR37 family may have evolved to link the sensation of a certain chemical class to regulation of innate behaviors or internal state in a way that has remained crucial across species.

Direct Links between Sensation and Action in the Evolution of Single OR Genes

Adaptation through change in the number of OR genes is only possible if the olfactory nervous system has ways to employ the new receptor genes that emerge as particular subfamilies grow and diversify. Because natural odor blends activate only a subset of receptors [22], expressing newly duplicated and diversified subfamily members may provide a more nuanced neural representation of the regions of chemical space they sense. This could in turn allow finer resolution in discriminating the makeup or concentrations of the odors sensed by the ancestral receptors.

What this model does not explain is how an animal's innate interpretation of a given odor may evolve. Many vertebrates exhibit hardwired patterns of behavior that can be released by precise chemical stimuli, including even some monomolecular odorants outside the context of their natural odor blends (Figure 3) [58,63,76,77]. Animals instead interpret these signals, at least in part, through single, precisely- and sensitively-tuned olfactory receptors that tap into hardwired, behavior-evoking neural circuits. This pattern of receptor tuning and neural organization is prominent in insects, perhaps because their olfactory receptor repertoires are much smaller (and less redundant) than those of vertebrates. The compounds that elicit innate responses in both insects and vertebrates are essential for survival and reproduction — sex pheromones, attractive or aversive food odors and predator signatures — explaining why olfactory neurons have evolved to detect them with high sensitivity and to couple directly to neural circuits that drive instinctive behaviors (Figure 3) [78–81]. In mice, for instance, the exocrine-secreted peptides ESP1 and ESP22 act exclusively through V2Rp5 and an unknown V2R to promote female sexual receptivity (the lordosis posture) and to block male sexual activity toward juveniles below reproductive age, respectively [82,83]. Similarly, phenethylamine found in the urine of carnivores acts by binding to the receptor TAAR4 to promote avoidance behavior [84,85]. These individual receptors therefore play non-redundant roles in shaping odor perception and interpretation, suggesting that even in olfactory systems that use hundreds or thousands of receptors, single members of these gene families have evolved outsized ecological significance and privileged neural access to specific central circuits [84].

This model also predicts that, in animal lineages in which the role of olfaction in driving stereotyped reproductive and defensive behaviors has waned, there will be few cases where

individual receptor genes are experiencing strong selective pressure. This seems to be the case in humans, for whom there is almost no evidence for positive selection in the coding sequences of single OR genes since the divergence from chimpanzees about seven million years ago. It is therefore unlikely that particular OR genes have conferred a selective advantage during human evolution [86].

Nonetheless, genetic variation in single human ORs — under selection or not — can produce large differences in odor perception. Detection thresholds and (un)pleasantness ratings of many odors have been linked to common polymorphisms in and around human OR genes. These include a receptor coding mutation that alters sensitivity to and valence of the male hormones androstenedione and androstenedione [87], a mutation that accounts for nearly all observed variation in the perception of beta-ionone and dramatically alters preference for foods emitting this odor [88], and an OR cluster-linked polymorphism that correlates with the aversive soapiness of cilantro [89]. Furthermore, as in other mammals, gland secretions can in a laboratory setting bias sexual attraction and nipple-orienting behavior in adult and infant humans, respectively, raising the possibility that we employ unidentified pheromones [90,91]. If these signals follow the pattern of other social and suckling pheromones, they may include monomolecular odorants acting through high-affinity, specialist receptors [77,92]. Thus, even if they are not under strong selection, the observation that single human ORs influence odor perception supports the possibility that they contribute meaningfully to innate behavior through unknown mechanisms.

Segregated Olfactory Pathways as Substrates for Innate Olfactory Behaviors

Defined odorants can drive innate responses or preferences through a few receptors among hundreds or thousands, many of which may be active simultaneously during typical sensation of complex odor blends [93]. How, then, are particular cues coupled precisely to appropriate odor-driven behaviors? It appears that the olfactory system begins to sort odor information in the peripheral sense organs. Several hardwired, parallel pathways — from sensory neurons to subregions of the olfactory bulb to distinct central brain targets — arose early in the evolution of vertebrates, using distinct receptor families and subfamilies to convey different categories of chemical signal. Although zebrafish, unlike mice, have only a single olfactory epithelium, the primary sensory neurons that express ORs and TAARs are molecularly, morphologically and anatomically distinct from those that express amino acid-sensing V2Rs. These two cell types — ciliated and microvillar sensory neurons, respectively — are found in different layers of the olfactory epithelium and project to separate regions of the olfactory bulb, presaging the separation of mammalian ciliated neurons (in the main olfactory epithelium) and microvillar neurons (in the vomeronasal organ) and their projections to distinct main and accessory olfactory bulbs [33,34]. The zebrafish epithelium also contains a third cell type (crypt neurons) that expresses one of the few zebrafish V1R genes in most cells and projects to a unique glomerulus [29,32,94]. In addition, a closely related receptor has recently been found to detect a pheromone that releases egg-laying behavior, reminiscent of the reproductive behaviors regulated by mammalian




Figure 4. Peripheral organization of OR family expression across vertebrates.

Top: A horizontal section of a quarter of an olfactory rosette, the water-exposed pit-like olfactory organ of the zebrafish (*Danio rerio*). Sensory neurons are embedded in a main olfactory epithelium (main olfactory epithelium) with a water-filled lumen. Members of four of the five major GPCR OR families are expressed in different sensory neuron types: TAARs and Class I classical ORs (here denoting also fish ORs that belong to neither Class I nor Class II) are expressed in ciliated cells occupying the basal layers of the epithelium; V2Rs in microvillar cells in the middle layer; and at least one V1R in crypt cells of the upper layer. Axons of the three cell types project to different, spatially segregated, and stereotyped glomeruli of the zebrafish olfactory bulb. Middle: A coronal section of one half of the main olfactory epithelium and the vomeronasal organ (vomeronasal organ) of the western clawed frog (*Xenopus laevis*). An air-filled medial diverticulum (MD) houses neurons expressing Class II classical ORs and several V1Rs; a water-filled lateral diverticulum (LD), Class I classical ORs, several TAARs, and ancestral V2Rs; the water-filled (vomeronasal organ), newer members of the expanded V2R family. Bottom: A coronal section of the main olfactory epithelium and vomeronasal organ of the house mouse (*Mus musculus*). Receptor family expression is segregated (several exceptions not pictured), with Class I ORs and TAARs expressed in the dorsal main olfactory epithelium, Class II ORs in the ventral main olfactory epithelium (and some in the dorsal main olfactory epithelium, not pictured), V1Rs in the apical vomeronasal organ, V2Rs in the basal vomeronasal organ, and FPRs in both vomeronasal organ layers. Some sensory neurons in the ‘cul-de-sacs’ of the main olfactory epithelium express the receptor guanylate cyclase GC-D and multiple members of a non-GPCR family of four-pass transmembrane chemoreceptors, Ms4a.

V1Rs (Figure 3) [95]. This “ancestral” organization of the olfactory periphery therefore suggests that discrete subsystems and the receptors they express are hardwired to yield different behavioral responses to odor (Figure 4).

Consistent with this possibility, a major olfactory adaptation along the amphibian and terrestrial branches of the vertebrate tree includes the development of segregated neuroanatomical structures that express more recently expanded OR subfamilies. Unlike fish, amphibians have evolved a separate vomeronasal organ that detects attractive peptide sex pheromones (specific to amphibians) through unusually large and recently expanded subfamilies of V2Rs, as well as an air-filled chamber of the main olfactory epithelium that expresses the massively enlarged Class II OR repertoire described above. Evolutionarily more ancient V2Rs tuned to single amino acids remain expressed in the water-filled main olfactory epithelium compartment along with V1Rs, Class I ORs, and TAARs (Figure 4) [62,96,97]. The observation that newer V2R genes have acquired both a novel expression pattern and a novel role in innate reproductive behaviors suggests that odor interpretation depends not only on what ligands are detected but also on the developmental identities and specialized sensory pathways of the cells that detect them [98].

The segregation of the olfactory periphery has proceeded even further in rodents. Mice have separate epithelial layers for V1Rs and V2Rs within the vomeronasal organ and distinct dorsoventral zones and molecular machinery for the expression of TAARs, Class I ORs, and Class II ORs in the main olfactory epithelium (Figure 4) [99]. Additional, smaller sensory structures include a group of neurons at the tip of the nose, the Grueneberg ganglion, which is thought to detect alarm pheromones and structurally related predator odorants [100,101], and a patch of olfactory neurons on the nasal septum that largely express a single broadly-tuned OR and may convey information about breathing [102,103]. Finally, within cul-de-sacs of the main olfactory epithelium resides a molecularly atypical set of neurons which each express both the receptor guanylate cyclase GC-D and a recently-described family of non-GPCR olfactory receptors (encoded by the *Membrane Spanning, 4-pass A (Ms4a)* genes); these cells are thought to mediate the social acquisition of food preference (Figure 4 bottom) [104–106].

The olfactory bulb appears to organize this variety of incoming sensory information into domains larger than a single glomerulus. For instance, the sensory neurons that express class I ORs, class II ORs, TAARs, or the MS4As/GC-D are developmentally constrained to innervate circumscribed and non-intersecting regions of the olfactory bulb (Figure 5) [99,107]. Furthermore, there is substantial evidence that activity in spatially and molecularly segregated glomeruli drives different patterns of stereotyped behavior via hardwired projections to central brain regions. First, an ‘innate aversion’ domain for predator odor sensation can be found at the ventral subdivision of the dorsal olfactory bulb, while the dorsalmost Class I-expressing domain is instead required for aversive responses to spoiled food odorants [108]. Second, the ventral olfactory bulb houses the glomeruli of molecularly distinct neurons (expressing components of an alternative, TrpM5-dependent chemotransduction pathway) that preferentially respond to urine-derived pheromones and project to the medial ‘vomeronasal amygdala’, as well as the cluster of OR37 glomeruli and their unusual mitral cell projections to the hypothalamus [75,77,109,110]. Third, the TAAR domain of the olfactory bulb is required for innate behavioral responses to several ecologically important amines, even though these same




Figure 5. Spatial and molecular organization of projection targets and behavioral responses downstream of distinct mouse olfactory bulb glomeruli.

Circumscribed zones of glomeruli are innervated by sensory neurons expressing each of the four largest GPCR OR families or the MS4As. In addition, Class I and Class II OR glomeruli occupy the dorsal-most and more ventral zones, respectively; and projections from Grueneberg ganglion sensory neurons form their own ‘necklace’ (light blue) anterior to the GC-D/MS4a ‘necklace’ (green). Dorsal Class II OR glomeruli form a molecularly defined zone, which includes glomeruli necessary for innate aversion of some predator odorants. Within the ventral zone of Class II OR glomeruli, some regions are enriched for the glomeruli of TrpM5- or OR37 subfamily-expressing sensory neurons.

odorants can be detected by receptors in the main olfactory system (Figure 5) [84]. Recent data also suggest a global organization to the main olfactory bulb, in which glomeruli that reside more rostrally evoke investigatory behaviors, while those that are more caudal evoke aversion [111]. Characterizing the mitral cell projections downstream of specific glomeruli may reveal different capacities to influence a variety of central processing or behavioral centers. This ‘switchboard’ organization could allow each OR gene to adapt its chemosensory tuning according to its domain’s function without altering the circuits downstream of the receptor.

However, the interpretation of an odor may not be determined simply by whether it recruits neural activity within a specific glomerular domain. Within the TAAR domain, for instance, the TAAR4 glomerulus is required for aversion to phenethylamine, but the nearby TAAR5 glomerulus is highly tuned to the attractive mouse pheromone trimethylamine [84,112]. Thus, even closely related receptors and their closely apposed glomeruli — within a similar subdomain of the olfactory bulb — may drive opposite behaviors. Combinatorial effects of multiple (TAAR and non-TAAR) glomeruli may produce the different behavioral responses to high-affinity TAAR4 and TAAR5 ligands. Consistent with this

possibility, the effect of TAAR5 activation appears to be inverted in rats, for which trimethylamine is aversive (Figure 3) [112]. However, it is also possible that, within the TAAR domain of the bulb, TAAR4 and TAAR5 glomeruli are connected to distinct downstream circuits that drive fundamentally different behaviors. Determining whether the differential effects of TAAR4 and TAAR5 ligands in mice rely upon glomerulus-specific projections to higher brain regions, the coordinated activity of groups of glomeruli, or on circuit activity within the TAAR domain will therefore reveal key principles of how odor representations in the olfactory bulb govern perception and behavior.

The appeal of the ‘domain hypothesis’ for coupling odors to behavior lies in its relative developmental simplicity. In this model, the olfactory system could take advantage of developmentally-specific gradients of morphogenetic and axon guidance molecules to build subcircuits within the olfactory bulb that connect to distinct higher brain regions. However, it is less clear how the olfactory system could differentially connect individual glomeruli within a given domain to specific higher brain circuits that mediate particular behaviors. One highly speculative possibility is that the olfactory bulb’s output layer of mitral cells is, like the retinal ganglion cell layer of the retina, tiled with molecularly distinct cell types that convey different types of information via their projection or spiking patterns. Hard-wired synaptic connections to these mitral cell types might then be biased by the complement of cell surface proteins on OSN axons, which are themselves determined by both their developmental identity and the OR gene stochastically chosen for expression [113]. This sort of coupling between ORs and central circuits could explain why many odors appear to have a stereotyped attractive or aversive valence in mice and humans, even when the ORs responsible for their detection are evolving neutrally [114,115]. In other words, behavioral biases could be substrates for adaptation (e.g. through changes in the expression of cell surface proteins given a particular OR choice) but would also necessarily exist by chance. Thus, the multiglomerular domains of the olfactory bulb may determine which downstream targets are available (e.g. medial amygdala from the accessory bulb and cortical amygdala from the main bulb) but leave enough flexibility for receptor-specific behaviors to evolve. The loss of several subsystems in humans may explain why single OR genes are not under strong selective pressure in our lineage, as a highly-tuned receptor would have few or no specialized circuits to tap into to generate innate behavioral responses.

Conclusion

How evolution sculpts the olfactory system is a response to both the general problem of probing a wide swath of chemical space, as well as the unique sensory challenges faced by each animal species. In vertebrates, both dramatic and subtle adaptations to the chemical environment are possible because of the vast, hierarchical structure of the OR gene families. First, rapid growth or pruning of large OR families accommodates major changes in habitat and lifestyle, as with ocean-to-terrestrial shifts and derived primacy of the visual sense. Second, selection on individual OR families or subfamilies permits niche adaptation or conserved chemical communication between conspecifics. Third, genetic variation in single OR genes, among thousands, can alter behaviors driven by precise chemical cues. The molecular logic and

neuroanatomical architecture of the vertebrate olfactory nervous system may have favored this dramatic pattern of olfactory receptor evolution, as newly arising olfactory receptor genes integrate by default into segregated pathways that influence both innate and learned behaviors. Peering into this dynamic process reveals a direct link between the evolution of an animal's genome and the tasks solved by its nervous system.

ACKNOWLEDGEMENTS

We thank Hannah Somhegyi for help designing figures and Taralyn Tan for contributing figure illustrations. D.M.B. is supported by a National Science Foundation predoctoral fellowship and a Sackler Scholarship in Psychobiology. J.M.L. is supported by the European Molecular Biology Organization (ALTF 379-2011), the Human Frontiers Science Program (LT001086/2012), and the Belgian American Educational Foundation. H.E.H. is an investigator of the Howard Hughes Medical Institute. S.R.D. is supported by fellowships from the Burroughs Wellcome Fund, the Vallee Foundation, and by grant RO11DC011558 from the National Institutes of Health.

REFERENCES

- Wyatt, T.D. (2014). Introduction to chemical signaling in vertebrates and invertebrates. In *Neurobiology of Chemical Communication*, C. Mucignat-Caretta, ed. (FL: Boca Raton).
- Meister, M. (2015). On the dimensionality of odor space. *Elife* 4, e07865.
- Bargmann, C.I. (2006). Comparative chemosensation from receptors to ecology. *Nature* 444, 295–301.
- Hansson, B.S., and Stensmyr, M.C. (2011). Evolution of insect olfaction. *Neuron* 72, 698–711.
- Zhang, X., and Firestein, S. (2002). The olfactory receptor gene superfamily of the mouse. *Nat. Neurosci.* 5, 124–133.
- Touhara, K., and Vosshall, L.B. (2009). Sensing odorants and pheromones with chemosensory receptors. *Annu. Rev. Physiol.* 71, 307–332.
- Finlay, B.L., and Darlington, R.B. (1995). Linked regularities in the development and evolution of mammalian brains. *Science* 268, 1578–1584.
- Meisami, E., and Bhatnagar, K.P. (1998). Structure and diversity in mammalian accessory olfactory bulb. *Microsc. Res. Tech.* 43, 476–499.
- Yopak, K.E., Lisney, T.J., and Collin, S.P. (2015). Not all sharks are “swimming noses”: variation in olfactory bulb size in cartilaginous fishes. *Brain Struct. Funct.* 220, 1127–1143.
- Manzini, I., and Korsching, S. (2011). The peripheral olfactory system of vertebrates: molecular, structural and functional basics of the sense of smell. *e-Neuroforum* 2, 68–77.
- Dalton, R.P., and Lomvardas, S. (2015). Chemosensory receptor specificity and regulation. *Annu. Rev. Neurosci.* 38, 331–349.
- Buck, L., and Axel, R. (1991). A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* 65, 175–187.
- Ressler, K.J., Sullivan, S.L., and Buck, L.B. (1993). A zonal organization of odorant receptor gene expression in the olfactory epithelium. *Cell* 73, 597–609.
- Johnson, M.A., Tsai, L., Roy, D.S., Valenzuela, D.H., Mosley, C., Magklara, A., Lomvardas, S., Liberles, S.D., and Barnea, G. (2012). Neurons expressing trace amine-associated receptors project to discrete glomeruli and constitute an olfactory subsystem. *Proc. Natl. Acad. Sci. USA* 109, 13410–13415.
- Dulac, C., and Axel, R. (1995). A novel family of genes encoding putative pheromone receptors in mammals. *Cell* 83, 195–206.
- Herrada, G., and Dulac, C. (1997). A novel family of putative pheromone receptors in mammals with a topographically organized and sexually dimorphic distribution. *Cell* 90, 763–773.
- Ryba, N.J., and Tirindelli, R. (1997). A new multigene family of putative pheromone receptors. *Neuron* 19, 371–379.
- Silvotti, L., Moiani, A., Gatti, R., and Tirindelli, R. (2007). Combinatorial co-expression of pheromone receptors, V2Rs. *J. Neurochem.* 103, 1753–1763.
- Riviere, S., Challet, L., Fluegge, D., Spehr, M., and Rodriguez, I. (2009). Formyl peptide receptor-like proteins are a novel family of vomeronasal chemosensory receptors. *Nature* 459, 574–577.
- Liberles, S.D., Horowitz, L.F., Kuang, D., Contos, J.J., Wilson, K.L., Siltberg-Liberles, J., Liberles, D.A., and Buck, L.B. (2009). Formyl peptide receptors are candidate chemosensory receptors in the vomeronasal organ. *Proc. Natl. Acad. Sci. USA* 106, 9842–9847.
- Mombaerts, P. (2004). Genes and ligands for odorant, vomeronasal and taste receptors. *Nat. Rev. Neurosci.* 5, 263–278.
- Lin da, Y., Shea, S.D., and Katz, L.C. (2006). Representation of natural stimuli in the rodent main olfactory bulb. *Neuron* 50, 937–949.
- Sosulski, D.L., Bloom, M.L., Cutforth, T., Axel, R., and Datta, S.R. (2011). Distinct representations of olfactory information in different cortical centres. *Nature* 472, 213–216.
- Choi, G.B., Stettler, D.D., Kallman, B.R., Bhaskar, S.T., Fleischmann, A., and Axel, R. (2011). Driving opposing behaviors with ensembles of piriform neurons. *Cell* 146, 1004–1015.
- Root, C.M., Denny, C.A., Hen, R., and Axel, R. (2014). The participation of cortical amygdala in innate, odour-driven behaviour. *Nature* 515, 269–273.
- Mohedano-Moriano, A., Pro-Sistiaga, P., Ubeda-Banon, I., Crespo, C., Insausti, R., and Martinez-Marcos, A. (2007). Segregated pathways to the vomeronasal amygdala: differential projections from the anterior and posterior divisions of the accessory olfactory bulb. *Eur. J. Neurosci.* 25, 2065–2080.
- Hussain, A., Saraiva, L.R., and Korsching, S.I. (2009). Positive Darwinian selection and the birth of an olfactory receptor clade in teleosts. *Proc. Natl. Acad. Sci. USA* 106, 4313–4318.
- Saraiva, L.R., Ahuja, G., Ivandic, I., Syed, A.S., Marioni, J.C., Korsching, S.I., and Logan, D.W. (2015). Molecular and neuronal homology between the olfactory systems of zebrafish and mouse. *Sci. Rep.* 5, 11487.
- Saraiva, L.R., and Korsching, S.I. (2007). A novel olfactory receptor gene family in teleost fish. *Genome. Res.* 17, 1448–1457.
- Weth, F., Nadler, W., and Korsching, S. (1996). Nested expression domains for odorant receptors in zebrafish olfactory epithelium. *Proc. Natl. Acad. Sci. USA* 93, 13321–13326.
- Sato, Y., Miyasaka, N., and Yoshihara, Y. (2007). Hierarchical regulation of odorant receptor gene choice and subsequent axonal projection of olfactory sensory neurons in zebrafish. *J. Neurosci.* 27, 1606–1615.
- Oka, Y., Saraiva, L.R., and Korsching, S.I. (2012). Crypt neurons express a single V1R-related ora gene. *Chem. Senses.* 37, 219–227.
- Sato, Y., Miyasaka, N., and Yoshihara, Y. (2005). Mutually exclusive glomerular innervation by two distinct types of olfactory sensory neurons revealed in transgenic zebrafish. *J. Neurosci.* 25, 4889–4897.
- Braubach, O.R., Fine, A., and Croll, R.P. (2012). Distribution and functional organization of glomeruli in the olfactory bulbs of zebrafish (*Danio rerio*). *J. Comp. Neurol.* 520, 2317–2339, Spc2311.
- Saito, H., Chi, Q., Zhuang, H., Matsunami, H., and Mainland, J.D. (2009). Odor coding by a Mammalian receptor repertoire. *Sci. Signal* 2, ra9.
- Nei, M., and Rooney, A.P. (2005). Concerted and birth-and-death evolution of multigene families. *Annu. Rev. Genet.* 39, 121–152.
- Zhang, X., and Firestein, S. (2009). Genomics of olfactory receptors. *Results. Probl. Cell. Differ.* 47, 25–36.
- Niimura, Y., Matsui, A., and Touhara, K. (2014). Extreme expansion of the olfactory receptor gene repertoire in African elephants and evolutionary dynamics of orthologous gene groups in 13 placental mammals. *Genome. Res.* 24, 1485–1496.

39. Sharon, D., Glusman, G., Pilpel, Y., Khen, M., Gruetzner, F., Haaf, T., and Lancet, D. (1999). Primate evolution of an olfactory receptor cluster: diversification by gene conversion and recent emergence of pseudogenes. *Genomics* 61, 24–36.
40. Neuhaus, E.M., Zhang, W., Gelis, L., Deng, Y., Noldus, J., and Hatt, H. (2009). Activation of an olfactory receptor inhibits proliferation of prostate cancer cells. *J. Biol. Chem.* 284, 16218–16225.
41. Niimura, Y., and Nei, M. (2006). Evolutionary dynamics of olfactory and other chemosensory receptor genes in vertebrates. *J. Hum. Genet.* 51, 505–517.
42. Adipietro, K.A., Mainland, J.D., and Matsunami, H. (2012). Functional evolution of mammalian odorant receptors. *PLoS Genet* 8, e1002821.
43. Liman, E.R. (2012). Changing senses: chemosensory signaling and primate evolution. *Adv. Exp. Med. Biol.* 739, 206–217.
44. Liman, E.R., and Innan, H. (2003). Relaxed selective pressure on an essential component of pheromone transduction in primate evolution. *Proc. Natl. Acad. Sci. USA* 100, 3328–3332.
45. Gilad, Y., Przeworski, M., and Lancet, D. (2004). Loss of olfactory receptor genes coincides with the acquisition of full trichromatic vision in primates. *PLoS Biol.* 2, E5.
46. Lucas, P.W., Dominy, N.J., Riba-Hernandez, P., Stoner, K.E., Yamashita, N., Loria-Calderon, E., Petersen-Pereira, W., Rojas-Duran, Y., Salas-Pena, R., Solis-Madrigal, S., *et al.* (2003). Evolution and function of routine trichromatic vision in primates. *Evolution* 57, 2636–2643.
47. Dominy, N.J., and Lucas, P.W. (2001). Ecological importance of trichromatic vision to primates. *Nature* 410, 363–366.
48. Matsui, A., Go, Y., and Niimura, Y. (2010). Degeneration of olfactory receptor gene repertoires in primates: no direct link to full trichromatic vision. *Mol. Biol. Evol.* 27, 1192–1200.
49. Zhao, H., Xu, D., Zhang, S., and Zhang, J. (2011). Widespread losses of vomeronasal signal transduction in bats. *Mol. Biol. Evol.* 28, 7–12.
50. Niimura, Y., and Nei, M. (2007). Extensive gains and losses of olfactory receptor genes in mammalian evolution. *PLoS One* 2, e708.
51. Niimura, Y., and Nei, M. (2005). Evolutionary dynamics of olfactory receptor genes in fishes and tetrapods. *Proc. Natl. Acad. Sci. USA* 102, 6039–6044.
52. Hayden, S., Bekaert, M., Crider, T.A., Mariani, S., Murphy, W.J., and Teeling, E.C. (2010). Ecological adaptation determines functional mammalian olfactory subgenomes. *Genome Res.* 20, 1–9.
53. Niimura, Y. (2009). Evolutionary dynamics of olfactory receptor genes in chordates: interaction between environments and genomic contents. *Hum. Genomics* 4, 107–118.
54. Freitag, J., Ludwig, G., Andreini, I., Rossler, P., and Breer, H. (1998). Olfactory receptors in aquatic and terrestrial vertebrates. *J. Comp. Physiol. A* 183, 635–650.
55. Freitag, J., Krieger, J., Strotmann, J., and Breer, H. (1995). Two classes of olfactory receptors in *Xenopus laevis*. *Neuron* 15, 1383–1392.
56. Mezler, M., Fleischer, J., and Breer, H. (2001). Characteristic features and ligand specificity of the two olfactory receptor classes from *Xenopus laevis*. *J. Exp. Biol.* 204, 2987–2997.
57. Li, Q., Tachie-Baffour, Y., Liu, Z., Baldwin, M.W., Kruse, A.C., and Liberles, S.D. (2015). Non-classical amine recognition evolved in a large clade of olfactory receptors. *Elife* 4, e10441.
58. Hussain, A., Saraiva, L.R., Ferrero, D.M., Ahuja, G., Krishna, V.S., Liberles, S.D., and Korsching, S.I. (2013). High-affinity olfactory receptor for the death-associated odor cadaverine. *Proc. Natl. Acad. Sci. USA* 110, 19579–19584.
59. Liberles, S.D. (2015). Trace amine-associated receptors: ligands, neural circuits, and behaviors. *Curr. Opin. Neurobiol.* 34, 1–7.
60. Rolen, S.H., Sorensen, P.W., Mattson, D., and Caprio, J. (2003). Polyamines as olfactory stimuli in the goldfish *Carassius auratus*. *J. Exp. Biol.* 206, 1683–1696.
61. Nei, M., Niimura, Y., and Nozawa, M. (2008). The evolution of animal chemosensory receptor gene repertoires: roles of chance and necessity. *Nat. Rev. Genet.* 9, 951–963.
62. Syed, A.S., Sansone, A., Nadler, W., Manzini, I., and Korsching, S.I. (2013). Ancestral amphibian v2rs are expressed in the main olfactory epithelium. *Proc. Natl. Acad. Sci. USA* 110, 7714–7719.
63. Kikuyama, S., Yamamoto, K., Iwata, T., and Toyoda, F. (2002). Peptide and protein pheromones in amphibians. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol.* 132, 69–74.
64. Grus, W.E., Shi, P., and Zhang, J. (2007). Largest vertebrate vomeronasal type 1 receptor gene repertoire in the semiaquatic platypus. *Mol. Biol. Evol.* 24, 2153–2157.
65. Isogai, Y., Si, S., Pont-Lezica, L., Tan, T., Kapoor, V., Murthy, V.N., and Dulac, C. (2011). Molecular organization of vomeronasal chemoreception. *Nature* 478, 241–245.
66. Hayden, S., Bekaert, M., Goodbla, A., Murphy, W.J., Davalos, L.M., and Teeling, E.C. (2014). A cluster of olfactory receptor genes linked to frugivory in bats. *Mol. Biol. Evol.* 31, 917–927.
67. Khan, I., Yang, Z., Maldonado, E., Li, C., Zhang, G., Gilbert, M.T., Jarvis, E.D., O'Brien, S.J., Johnson, W.E., and Antunes, A. (2015). Olfactory receptor subgenomes linked with broad ecological adaptations in Saurapsida. *Mol. Biol. Evol.* 32, 2832–2843.
68. Yang, H., Shi, P., Zhang, Y.P., and Zhang, J. (2005). Composition and evolution of the V2r vomeronasal receptor gene repertoire in mice and rats. *Genomics* 86, 306–315.
69. Papes, F., Logan, D.W., and Stowers, L. (2010). The vomeronasal organ mediates interspecies defensive behaviors through detection of protein pheromone homologs. *Cell* 141, 692–703.
70. Chameró, P., Marton, T.F., Logan, D.W., Flanagan, K., Cruz, J.R., Saghatelian, A., Cravatt, B.F., and Stowers, L. (2007). Identification of protein pheromones that promote aggressive behaviour. *Nature* 450, 899–902.
71. Hoppe, R., Lambert, T.D., Samollow, P.B., Breer, H., and Strotmann, J. (2006). Evolution of the “OR37” subfamily of olfactory receptors: a cross-species comparison. *J. Mol. Evol.* 62, 460–472.
72. Hoppe, R., Breer, H., and Strotmann, J. (2003). Organization and evolutionary relatedness of OR37 olfactory receptor genes in mouse and human. *Genomics* 82, 355–364.
73. Bautze, V., Bar, R., Fissler, B., Trapp, M., Schmidt, D., Beifuss, U., Bufe, B., Zufall, F., Breer, H., and Strotmann, J. (2012). Mammalian-specific OR37 receptors are differentially activated by distinct odorous fatty aldehydes. *Chem. Senses.* 37, 479–493.
74. Strotmann, J., Conzelmann, S., Beck, A., Feinstein, P., Breer, H., and Mombaerts, P. (2000). Local permutations in the glomerular array of the mouse olfactory bulb. *J. Neurosci.* 20, 6927–6938.
75. Bader, A., Klein, B., Breer, H., and Strotmann, J. (2012). Connectivity from OR37 expressing olfactory sensory neurons to distinct cell types in the hypothalamus. *Front Neural Circuits* 6, 84.
76. Dulac, C., and Torello, A.T. (2003). Molecular detection of pheromone signals in mammals: from genes to behaviour. *Nat. Rev. Neurosci.* 4, 551–562.
77. Lin, D.Y., Zhang, S.Z., Block, E., and Katz, L.C. (2005). Encoding social signals in the mouse main olfactory bulb. *Nature* 434, 470–477.
78. Dekker, T., Ibba, I., Siju, K.P., Stensmyr, M.C., and Hansson, B.S. (2006). Olfactory shifts parallel superspecialism for toxic fruit in *Drosophila melanogaster* sibling, *D. sechellia*. *Curr. Biol.* 16, 101–109.
79. Kurtovic, A., Widmer, A., and Dickson, B.J. (2007). A single class of olfactory neurons mediates behavioural responses to a *Drosophila* sex pheromone. *Nature* 446, 542–546.
80. Sakurai, T., Mitsuno, H., Mikami, A., Uchino, K., Tabuchi, M., Zhang, F., Sezutsu, H., and Kanzaki, R. (2015). Targeted disruption of a single sex pheromone receptor gene completely abolishes *in vivo* pheromone response in the silkworm. *Sci. Rep.* 5, 11001.

81. Stensmyr, M.C., Dweck, H.K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J., Grabe, V., Steck, K., Lavista-Llanos, S., *et al.* (2012). A conserved dedicated olfactory circuit for detecting harmful microbes in *Drosophila*. *Cell* *151*, 1345–1357.
82. Haga, S., Hattori, T., Sato, T., Sato, K., Matsuda, S., Kobayakawa, R., Sakano, H., Yoshihara, Y., Kikusui, T., and Touhara, K. (2010). The male mouse pheromone ESP1 enhances female sexual receptive behaviour through a specific vomeronasal receptor. *Nature* *466*, 118–122.
83. Ferrero, D.M., Moeller, L.M., Osakada, T., Horio, N., Li, Q., Roy, D.S., Cichy, A., Spehr, M., Touhara, K., and Liberles, S.D. (2013). A juvenile mouse pheromone inhibits sexual behaviour through the vomeronasal system. *Nature* *502*, 368–371.
84. Dewan, A., Pacifico, R., Zhan, R., Rinberg, D., and Bozza, T. (2013). Non-redundant coding of aversive odours in the main olfactory pathway. *Nature* *497*, 486–489.
85. Ferrero, D.M., Lemon, J.K., Fluegge, D., Pashkovski, S.L., Korzan, W.J., Datta, S.R., Spehr, M., Fendt, M., and Liberles, S.D. (2011). Detection and avoidance of a carnivore odor by prey. *Proc. Natl. Acad. Sci. USA* *108*, 11235–11240.
86. Gimelbrant, A.A., Skaletsky, H., and Chess, A. (2004). Selective pressures on the olfactory receptor repertoire since the human-chimpanzee divergence. *Proc. Natl. Acad. Sci. USA* *101*, 9019–9022.
87. Keller, A., Zhuang, H., Chi, Q., Vosshall, L.B., and Matsunami, H. (2007). Genetic variation in a human odorant receptor alters odour perception. *Nature* *449*, 468–472.
88. Jaeger, S.R., McRae, J.F., Bava, C.M., Beresford, M.K., Hunter, D., Jia, Y., Chheang, S.L., Jin, D., Peng, M., Gamble, J.C., *et al.* (2013). A Mendelian trait for olfactory sensitivity affects odor experience and food selection. *Curr. Biol.* *23*, 1601–1605.
89. Eriksson, N., Wu, S., Do, C.B., Kiefer, A.K., Tung, J.Y., Mountain, J.L., Hinds, D.A., and Francke, U. (2012). A genetic variant near olfactory receptor genes influences cilantro preference. *Flavour* *1*, 1.
90. Gelstein, S., Yeshurun, Y., Rozenkrantz, L., Shushan, S., Frumin, I., Roth, Y., and Sobel, N. (2011). Human tears contain a chemosignal. *Science* *331*, 226–230.
91. Varendi, H., and Porter, R.H. (2001). Breast odour as the only maternal stimulus elicits crawling towards the odour source. *Acta. Paediatr.* *90*, 372–375.
92. Schaal, B., Coureaud, G., Langlois, D., Ginies, C., Semon, E., and Perrier, G. (2003). Chemical and behavioural characterization of the rabbit mammary pheromone. *Nature* *424*, 68–72.
93. Rokni, D., Hemmelder, V., Kapoor, V., and Murthy, V.N. (2014). An olfactory cocktail party: figure-ground segregation of odorants in rodents. *Nat. Neurosci.* *17*, 1225–1232.
94. Ahuja, G., Ivandic, I., Salturk, M., Oka, Y., Nadler, W., and Korsching, S.I. (2013). Zebrafish crypt neurons project to a single, identified mediadorsal glomerulus. *Sci. Rep.* *3*, 2063.
95. Behrens, M., Frank, O., Rawel, H., Ahuja, G., Potting, C., Hofmann, T., Meyerhof, W., and Korsching, S. (2014). ORA1, a zebrafish olfactory receptor ancestral to all mammalian V1R genes, recognizes 4-hydroxyphenylacetic acid, a putative reproductive pheromone. *J. Biol. Chem.* *289*, 19778–19788.
96. Syed, A.S., Sansone, A., Roner, S., Bozorg Nia, S., Manzini, I., and Korsching, S.I. (2015). Different expression domains for two closely related amphibian TAARs generate a bimodal distribution similar to neuronal responses to amine odors. *Sci. Rep.* *5*, 13935.
97. Date-Ito, A., Ohara, H., Ichikawa, M., Mori, Y., and Hagino-Yamagishi, K. (2008). *Xenopus* V1R vomeronasal receptor family is expressed in the main olfactory system. *Chem. Senses.* *33*, 339–346.
98. Gliem, S., Syed, A.S., Sansone, A., Kludt, E., Tantalaki, E., Hassenklover, T., Korsching, S.I., and Manzini, I. (2013). Bimodal processing of olfactory information in an amphibian nose: odor responses segregate into a medial and a lateral stream. *Cell. Mol. Life. Sci.* *70*, 1965–1984.
99. Bozza, T., Vassalli, A., Fuss, S., Zhang, J.J., Weiland, B., Pacifico, R., Feinstein, P., and Mombaerts, P. (2009). Mapping of class I and class II odorant receptors to glomerular domains by two distinct types of olfactory sensory neurons in the mouse. *Neuron* *61*, 220–233.
100. Brechbuhl, J., Moine, F., Klaey, M., Nenniger-Tosato, M., Humi, N., Sporkert, F., Giroud, C., and Broillet, M.C. (2013). Mouse alarm pheromone shares structural similarity with predator scents. *Proc. Natl. Acad. Sci. USA* *110*, 4762–4767.
101. Brechbuhl, J., Klaey, M., and Broillet, M.C. (2008). Grueneberg ganglion cells mediate alarm pheromone detection in mice. *Science* *321*, 1092–1095.
102. Grosmaître, X., Fuss, S.H., Lee, A.C., Adipietro, K.A., Matsunami, H., Mombaerts, P., and Ma, M. (2009). SR1, a mouse odorant receptor with an unusually broad response profile. *J. Neurosci.* *29*, 14545–14552.
103. Grosmaître, X., Santarelli, L.C., Tan, J., Luo, M., and Ma, M. (2007). Dual functions of mammalian olfactory sensory neurons as odor detectors and mechanical sensors. *Nat. Neurosci.* *10*, 348–354.
104. Greer, P.L., Bear, D.M., Lassance, J.M., Bloom, M.L., Tsukahara, T., Pashkovski, S.L., Masuda, F.K., Nowlan, A.C., Kirchner, R., Hoekstra, H.E., *et al.* (2016). A family of non-GPCR chemosensors defines an alternative logic for mammalian olfaction. *Cell* *165*, 1734–1748.
105. Munger, S.D., Leinders-Zufall, T., McDougall, L.M., Cockerham, R.E., Schmid, A., Wandernoth, P., Wennemuth, G., Biel, M., Zufall, F., and Kelliher, K.R. (2010). An olfactory subsystem that detects carbon disulfide and mediates food-related social learning. *Curr. Biol.* *20*, 1438–1444.
106. Hu, J., Zhong, C., Ding, C., Chi, Q., Walz, A., Mombaerts, P., Matsunami, H., and Luo, M. (2007). Detection of near-atmospheric concentrations of CO₂ by an olfactory subsystem in the mouse. *Science* *317*, 953–957.
107. Pacifico, R., Dewan, A., Cawley, D., Guo, C., and Bozza, T. (2012). An olfactory subsystem that mediates high-sensitivity detection of volatile amines. *Cell Rep.* *2*, 76–88.
108. Kobayakawa, K., Kobayakawa, R., Matsumoto, H., Oka, Y., Imai, T., Ikawa, M., Okabe, M., Ikeda, T., Itohara, S., Kikusui, T., *et al.* (2007). Innate versus learned odour processing in the mouse olfactory bulb. *Nature* *450*, 503–508.
109. Lin, W., Margolskee, R., Donnert, G., Hell, S.W., and Restrepo, D. (2007). Olfactory neurons expressing transient receptor potential channel M5 (TRPM5) are involved in sensing semiochemicals. *Proc. Natl. Acad. Sci. USA* *104*, 2471–2476.
110. Thompson, J.A., Salcedo, E., Restrepo, D., and Finger, T.E. (2012). Second-order input to the medial amygdala from olfactory sensory neurons expressing the transduction channel TRPM5. *J. Comp. Neurol.* *520*, 1819–1830.
111. Kermen, F., Midroit, M., Kuczewski, N., Forest, J., Thevenet, M., Sacquet, J., Benetollo, C., Richard, M., Didier, A., and Mandairon, N. (2016). Topographical representation of odor hedonics in the olfactory bulb. *Nat. Neurosci.* *19*, 876–878.
112. Li, Q., Korzan, W.J., Ferrero, D.M., Chang, R.B., Roy, D.S., Buchi, M., Lemon, J.K., Kaur, A.W., Stowers, L., Fendt, M., *et al.* (2013). Synchronous evolution of an odor biosynthesis pathway and behavioral response. *Curr. Biol.* *23*, 11–20.
113. Serizawa, S., Miyamichi, K., Takeuchi, H., Yamagishi, Y., Suzuki, M., and Sakano, H. (2006). A neuronal identity code for the odorant receptor-specific and activity-dependent axon sorting. *Cell* *127*, 1057–1069.
114. Saraiva, L.R., Kondoh, K., Ye, X., Yoon, K.H., Hernandez, M., and Buck, L.B. (2016). Combinatorial effects of odorants on mouse behavior. *Proc. Natl. Acad. Sci. USA* *113*, E3300–3306.
115. Haddad, R., Medhanie, A., Roth, Y., Harel, D., and Sobel, N. (2010). Predicting odor pleasantness with an electronic nose. *PLoS Comput. Biol.* *6*, e1000740.