

Interspecific variation in cooperative burrowing behavior by *Peromyscus* mice

Nicole L. Bedford,^{1,2,3,4}  Jesse N. Weber,^{2,3,5}  Wenfei Tong,^{2,3} Felix Baier,^{1,2,6}  Ariana Kam,³ Rebecca A. Greenberg,³ and Hopi E. Hoekstra^{1,2,3,6,7,8} 

¹Howard Hughes Medical Institute, Harvard University, Cambridge, Massachusetts 02138, USA

²Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138, USA

³Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts 02138, USA

⁴Current address: Department of Zoology and Physiology, University of Wyoming, Laramie, Wyoming 82071, USA

⁵Current address: Department of Integrative Biology, University of Wisconsin–Madison, Madison, Wisconsin 53706, USA

⁶Department of Molecular and Cellular Biology, Harvard University, Cambridge, Massachusetts 02138, USA

⁷Center for Brain Science, Harvard University, Cambridge, Massachusetts 02138, USA

⁸E-mail: hoekstra@oeb.harvard.edu

Received May 19, 2021

Accepted June 12, 2022

Animals often adjust their behavior according to social context, but the capacity for such behavioral flexibility can vary among species. Here, we test for interspecific variation in behavioral flexibility by comparing burrowing behavior across three species of deer mice (genus *Peromyscus*) with divergent social systems, ranging from promiscuous (*Peromyscus leucopus* and *Peromyscus maniculatus*) to monogamous (*Peromyscus polionotus*). First, we compared the burrows built by individual mice to those built by pairs of mice in all three species. Although burrow length did not differ in *P. leucopus* or *P. maniculatus*, we found that *P. polionotus* pairs cooperatively constructed burrows that were nearly twice as long as those built by individuals and that opposite-sex pairs dug longer burrows than same-sex pairs. Second, to directly observe cooperative digging behavior in *P. polionotus*, we designed a burrowing assay in which we could video-record active digging in narrow, transparent enclosures. Using this novel assay, we found, unexpectedly, that neither males nor females spent more time digging with an opposite-sex partner. Rather, we demonstrate that opposite-sex pairs are more socially cohesive and thus more efficient digging partners than same-sex pairs. Together, our study demonstrates how social context can modulate innate behavior and offers insight into how differences in behavioral flexibility may evolve among closely related species.

KEY WORDS: Animal architecture, burrow, cooperation, deer mouse, social cohesion.

Although some innate behaviors are invariantly expressed, others are labile, allowing animals to flexibly adjust their behavior according to context. Notably, this capacity for behavioral flexibility can vary among closely related species. For example, the same behavior may be fixed in one species but flexible in another (Kappeler et al. 2013; Royle et al. 2014). Such interspecific variation raises the possibility that behavioral flexibility itself can be heritable and therefore subject to natural selection. Thus, under-

standing the conditions under which selection may favor the fixed or flexible expression of innate behavior has been of longstanding interest in evolutionary biology (Mayr 1974).

One hypothesis is that differences in social system contribute to interspecific variation in the capacity for behavioral flexibility (Bond et al. 2007; Amici et al. 2008). Specifically, species that vary in social organization, mating system, and/or parental care strategy experience persistent differences in social environment that may promote or constrain the evolution of flexible behavior. Within species, how the social environment affects the expression

Nicole L. Bedford and Jesse N. Weber contributed equally to this work.

of innate (i.e., unlearned) behavior is increasingly well understood. For example, social context—particularly group size—is known to modulate innate behaviors, such as odor preference in flies (Ramdya et al. 2015), mating behavior in water striders (Montiglio et al. 2017), and exploratory behavior in zebrafish (Guayasamin et al. 2017). Similarly, past social experience can alter territoriality in flies (Hoffmann 1990; Watanabe et al. 2017), social competence in cichlids (Taborsky et al. 2012), aggression in mice (Nelson et al. 2013; Stagkourakis et al. 2020), and nest building in rats (Sharpe 1975). Thus, although social context can be an important driver of behavioral flexibility within species, how the capacity for behavioral flexibility varies among species remains less clear.

Comparisons of closely related species with different social systems may be especially informative for understanding the role of social environment (if any) in the evolution of behavioral flexibility. Here, we focused on the effect of social context on burrowing behavior in three closely related species of deer mice (genus *Peromyscus*) with divergent social systems. The oldfield mouse (*Peromyscus polionotus*) is both socially and genetically monogamous (Foltz 1981), provides biparental care (Bendesky et al. 2017), and commonly nests in opposite-sex pairs (Blair 1951). By contrast, its sister species, the deer mouse (*Peromyscus maniculatus*), and an outgroup species, the white-footed mouse (*Peromyscus leucopus*), are highly promiscuous (Xia and Millar 1991; Ribble and Millar 1996), provide uniparental care (Xia and Millar 1988; Bendesky et al. 2017), and most often nest solitarily (Wolff and Hurlbutt 1982). Importantly, all three species construct burrows that can be used to avoid predators, cache food, or rear young.

Across *Peromyscus*, burrow architecture is both species-specific and innate. For example, animals reared in captivity for several generations, without exposure to soil or opportunity to burrow, recapitulate their species-specific burrow architecture when tested in the lab (Weber and Hoekstra 2009; Weber et al. 2013). Moreover, interspecific cross-fostering shows that mice raised by another *Peromyscus* species dig burrows typical of their own species, rather than those of their foster parents (Metz et al. 2017). These data are consistent with a strong genetic basis for interspecific variation in *Peromyscus* burrowing behavior. In both the lab and the field, *P. polionotus* builds “complex” burrows consisting of a long entrance tunnel, nest chamber, and upward sloping escape tunnel, whereas *P. maniculatus* and *P. leucopus* build “simple” burrows with a short entrance tunnel and terminal nest chamber (Dawson et al. 1988; Weber and Hoekstra 2009; Weber et al. 2013). However, it is unknown whether these species-specific architectures are fixed, or if mice flexibly adjust their burrow length according to social context.

Here, under controlled laboratory conditions, we tested mice from three *Peromyscus* species to determine if social context (i.e.,

digging alone or in pairs) affects burrow length. Because social system differs among these species, we may expect the ability to flexibly alter burrow length according to social context to differ as well. Specifically, we predicted that the monogamous *P. polionotus* would adjust its burrowing behavior based on social context, whereas the promiscuous *P. maniculatus* and *P. leucopus* would not. We found that the ability to alter burrow length according to social context varies among species: burrow length was unchanged between individual and pair trials in *P. maniculatus* and *P. leucopus*, whereas *P. polionotus* pairs cooperatively constructed burrows that were nearly twice as long as those dug by individuals. Moreover, by directly measuring digging behavior in the cooperatively burrowing species (*P. polionotus*), we show that digging efficiency and burrow length depend on the sex of one’s digging partner. Together, these results underscore how social context can differentially affect the expression of an innate behavior, even among closely related species.

Methods

Please see the Supporting Information.

Results

EFFECT OF SOCIAL CONTEXT ON BURROW LENGTH

To investigate the effect of social context on burrowing behavior, we first compared the burrows built by individual mice to those built by pairs of mice in three *Peromyscus* species. To do so, we introduced either one or two mice into large, sand-filled enclosures. After two overnight periods, we removed the mice and made casts of the resultant burrows with polyurethane filling foam. For pair trials, we noted whether the two mice were found in the same or separate burrows at the end of the trial. We then measured each burrow cast to determine burrow length and shape. For each trial, we recorded the number of burrows, measured the length of each burrow, and categorized the shape of each burrow as either “simple” or “complex.” We further noted the length of the longest burrow produced each trial, or “maximum burrow length” (see *Methods* in the Supporting Information).

In both individual and pair trials, mice from all three species produced species-typical burrows. *Peromyscus leucopus* and *P. maniculatus* dug “simple” burrows with a short entrance tunnel and terminal nest chamber, whereas *P. polionotus* dug “complex” burrows with a long entrance tunnel, nest chamber, and escape tunnel (Fig. 1A). However, using a linear mixed-effects model (LMM) controlling for mouse ID and sex, we found that pairs dug more burrows than individuals (individuals: 1.15 ± 0.04 burrows, pairs: 1.54 ± 0.07 burrows) in all three species (Fig. S1A; LMM, species: $F = 2.38$,




Figure 1. *Peromyscus polionotus* pairs cooperatively construct long burrows. (A) Phylogenetic relationships, presence/absence of derived traits, and species-typical burrow architectures of three *Peromyscus* species. (B) Ratio of pair to individual burrow lengths in three species. A ratio > 1 indicates that pairs dug longer burrows than individuals. All pair-types (FF, MM, FM) and both sexes (F, M) are included in ratio calculations (*P. leucopus*: n = 9 pairs, 15 individuals; *P. maniculatus*: n = 22 pairs, 21 individuals; *P. polionotus*: n = 52 pairs, 52 individuals). (C) Maximum burrow length dug by individuals (F = purple, M = green) and pairs of mice (FF = dark purple, MM = dark green, FM = gold) in three species. Data are plotted on a square-root scale. Data points represent the mean of 1–7 trials per individual or unique pair. (D) Ratio of observed to expected burrow lengths in *P. polionotus* pairs, given the known output of individuals comprising the pair. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Error bars represent SEM.

$P = 0.098$; context: $F = 31.32$, $P < 0.001$; species $\times$ context: $F = 1.38$, $P = 0.253$). Nonetheless, two mice were found together in the same burrow in 58% of *P. leucopus* trials, 88% of *P. maniculatus* trials, and 100% of *P. polionotus* trials. In addition, maximum burrow length differed significantly among species and between trial-types (Fig. S1B; LMM, species: $F = 157.41$, $P < 0.001$; context: $F = 21.56$, $P < 0.001$; species $\times$ context: $F = 20.51$, $P < 0.001$). Using planned contrasts, we found an effect of social context on maximum burrow

length only in *P. polionotus* (Fig. 1B; *P. leucopus*: $t = 0.16$, $P = 0.871$; *P. maniculatus*: $t = 0.89$, $P = 0.374$; *P. polionotus*: $t = 11.27$, $P < 0.001$). *Peromyscus polionotus* pairs dug, on average, 82% longer burrows than *P. polionotus* individuals (individuals: 35.3 ± 2.1 cm, pairs: 64.3 ± 4.2 cm). By contrast, in *P. leucopus* and *P. maniculatus*, the maximum burrow length in individual and pair trials was statistically indistinguishable (*P. leucopus* individuals: 7.0 ± 1.2 cm, *P. leucopus* pairs: 6.9 ± 1.5 cm; *P. maniculatus* individuals: 15.0 ± 1.0 cm,

P. maniculatus pairs: 16.4 ± 0.9 cm). Thus, pairs of mice dug more burrows than individuals in all three species, but the effect of social context on burrow length varied by species: *P. polionotus* dug significantly longer burrows when digging in pairs, whereas its sister species (*P. maniculatus*) and an outgroup species (*P. leucopus*) did not.

Given that we observed social modulation of burrow length only in *P. polionotus*, we further investigated the effect of trial-type (i.e., female or male individuals; same-sex or opposite-sex pairs) on maximum burrow length in this species. Controlling for mouse ID, we found a significant effect of trial-type (Fig. 1C; LMM, $F = 28.52$, $P < 0.001$). Specifically, males dug longer burrows than females (planned contrasts, $t = 2.42$, $P = 0.017$) and opposite-sex pairs dug longer burrows than same-sex pairs ($t = 3.44$, $P = 0.001$). Male burrows were, on average, 54% longer than female burrows (male: 44.6 ± 2.7 cm, female: 29.0 ± 2.2 cm) and burrows dug by opposite-sex pairs were, on average, 28% longer than those dug by same-sex pairs (opposite-sex: 72.8 ± 6.4 cm, same-sex: 56.8 ± 3.8 cm).

To control for sex differences and individual variation in burrowing output, we tested whether the burrows dug by different pair-types (i.e., same-sex or opposite-sex) were shorter or longer than expected, given the typical burrow length dug by individuals comprising the pair. For each unique *P. polionotus* pair, we calculated the observed:expected burrow length ratio, where the observed value is the mean burrow length dug by a given pair, and the expected value is the sum of mean burrow lengths dug individually by each member of the pair. Controlling for mouse ID and partner ID, we found that opposite-sex pairs dug longer burrows than expected compared to same-sex pairs (Fig. 1D; LMM, $F = 9.12$, $P = 0.005$). The observed:expected burrow length ratio was 0.82 ± 0.05 for same-sex pairs and 1.18 ± 0.14 for opposite-sex pairs. This ratio was significantly less than 1 for same-sex pairs, but statistically indistinguishable from 1 for opposite-sex pairs (t -tests, FF: $t = 2.53$, $P = 0.012$; MM: $t = 2.97$, $P = 0.006$; FM: $t = 1.26$, $P = 0.234$), indicating that per capita output declines with a same-sex partner, but is unchanged with an opposite-sex partner. Together, these data suggest that *P. polionotus* mice alter their burrowing output according to social context (i.e., whether digging with a same- or opposite-sex partner).

EFFECT OF REPRODUCTIVE STATE ON FEMALE BURROW LENGTH

One potential explanation for the observation that opposite-sex pairs dig longer burrows than same-sex pairs is that females, when paired with a male, upregulate their digging behavior. Indeed, early studies of burrowing behavior in *P. polionotus* hypothesized that pregnancy may induce increased burrowing output in females (Dawson et al. 1988), as has been documented in other




Figure 2. *Peromyscus polionotus* females dig longer burrows after being co-housed with a male. (A) Experimental design schematic. Individual females were first tested as virgins to determine baseline burrowing output (top). After a period of cohabitation with a conspecific male (middle), females were tested again (bottom). By chance, 21/42 females became pregnant during the cohabitation period. (B) Maximum burrow length dug by females before (left) and after (right) male cohabitation. Each line represents the mean of two trials per individual, per timepoint. * $P < 0.05$. Error bars represent SEM.

species such as rabbits (Gonzalez-Mariscal et al. 2007). We therefore tested if *P. polionotus* females that were paired with a male dug longer burrows than those housed only with same-sex mice. We introduced single virgin females into large, sand-filled enclosures and, after one overnight period, removed the mouse and cast the resultant burrows. We then co-housed each female with a conspecific male and re-tested the female's individual burrowing behavior (Fig. 2A). By chance, half of the females (21/42) became pregnant during the cohabitation period. Notably, we found no effect of virgin burrow length on the probability of becoming pregnant (GLMM, $z = 0.21$, $P = 0.833$). Controlling for mouse ID, we found a significant effect of pairing with a male, but not pregnancy, on the length of the longest burrow dug per trial (Fig. 2B; LMM, cohabitation: $F = 5.26$, $P = 0.023$, pregnancy: $F = 1.03$, $P = 0.317$, cohabitation $\times$ pregnancy: $F = 0.01$, $P = 0.905$). *Peromyscus polionotus* females, regardless of pregnancy status, dug longer burrows after cohabitation with a male, with a median increase of 11% over their previous trials (virgin: 22.0 ± 1.4 cm, post-cohabitation: 26.2 ± 2.6 cm). This contrasts with *P. maniculatus* females, where we found a median burrow length increase of 21% in pregnant females only (Fig. S2, pregnant: $t = 3.01$, $P = 0.003$, non-pregnant: $t = 0.93$, $P = 0.355$).

We next asked whether this 11% increase in burrowing output by *P. polionotus* females that had been paired with a male could explain the observation that opposite-sex pairs dig

longer burrows than same-sex pairs (Fig. 1D). To account for this 11% increase, we adjusted the observed:expected burrow length ratio of opposite-sex pairs by increasing the expected burrow length value for individual females by 11%. However, even with this adjusted expected value, we still found significantly greater observed:expected burrow length ratios for opposite-sex versus same-sex pairs (LMM, $F = 7.01$, $P = 0.012$), suggesting that up-regulation of digging behavior by paired females alone does not fully explain the longer burrows produced by opposite-sex pairs.

NOVEL ASSAY TO MEASURE INDIVIDUAL BEHAVIOR

Although our large enclosures are semi-naturalistic arenas in which mice dig species-typical burrows, this behavioral assay does not allow us to directly observe digging or determine the relative contributions of individual mice during pair trials. We therefore designed a narrow, transparent sand-filled enclosure to directly observe their nocturnal, underground behavior (Fig. 3A). In brief, we introduced either one or two mice at the start of the dark cycle into narrow enclosures and recorded 8 hours of overnight video under infrared light and, for all pair trials, manually scored two 10-minute observation periods per trial (see *Methods* in the Supporting Information). For each observation period, we recorded all burrow entries and exits and specific digging behaviors for both mice (see *Video analysis*). We also measured burrow length using still images from the video taken at the beginning and end of each observation period as well as at the end of the 8-hour trial (see *Photo analysis*).

First, to first confirm that burrowing behavior in the narrow, transparent enclosures is similar to that observed in the large enclosures, we compared the length of the longest burrow dug per trial by individuals versus pairs of mice. As in the large enclosures, *P. polionotus* pairs dug more burrows (Fig. S3A; LMM controlling for mouse ID and sex, $F = 23.00$, $P < 0.001$) and longer burrows than *P. polionotus* individuals (Fig. S3B; LMM, $F = 23.29$, $P < 0.001$). We also found a significant effect of trial-type on maximum burrow length (Fig. S3C; LMM controlling for mouse ID, $F = 9.69$, $P < 0.001$). Specifically, males dug longer burrows than females (planned contrasts, $t = 2.27$, $P = 0.024$) and opposite-sex pairs dug longer burrows than same-sex pairs ($t = 2.45$, $P = 0.016$). Thus, the effect of social context on burrow length in *P. polionotus* was consistent between our two behavior arenas.

SEX DIFFERENCES IN BURROW LENGTH

In the transparent enclosures, we used a repeated-measures design to compare the burrowing behavior of 16 females and 16 males across same-sex and opposite-sex trials (Fig. 3B). We first tested all animals as individuals to determine baseline burrowing output and then tested individuals with a same-sex and opposite-sex partner, in random order. We found that total burrow length

differed significantly between sexes and among trial-types (LMM controlling for mouse ID, sex: $F = 5.19$, $P = 0.030$; context: $F = 27.56$, $P < 0.001$; sex $\times$ context: $F = 3.52$, $P = 0.032$). Compared to individual trials, total burrow length for females increased by 47% in same-sex trials and 87% in opposite-sex trials (Fig. 3C; Tukey contrasts, F vs. FF: $t = 3.09$, $P = 0.007$; F vs. FM: $t = 5.78$, $P < 0.001$; FF vs. FM: $t = 2.74$, $P = 0.019$). By contrast, total burrow length for males increased by 51% in pair trials, regardless of pair-type (Fig. 3C; Tukey contrasts, M vs. MM: $t = 3.29$, $P = 0.003$; M vs. FM: $t = 3.30$, $P = 0.003$; MM vs. FM: $t = 0.01$, $P = 0.999$). These results suggest that females and males may respond differently to changes in social context, but direct observation of individual-level behavior is necessary to address this question.

SEX DIFFERENCES IN BURROWING BEHAVIOR

To better understand how females and males behave when digging in pairs, we quantified individual-level behavior during pair trials. For each mouse in a pair, we recorded underground behavior, digging behavior, and hind-kicks over four 10-minute observation periods per mouse, per social context (Fig. S4). We first pooled data across pair-types to explore sex differences in behavior, using GLMMs with a Poisson link function to analyze number of bouts and LMMs to analyze mean bout duration and total duration (all models included mouse sex and pair-type as fixed effects and mouse ID, observer ID, trial number, and observation period as random effects). When controlling for social context, we found that males went underground 40% more often, dug 47% more often, and performed 113% more hind-kicks than females (Fig. 3D; GLMMs, underground: $z = 4.01$, $P < 0.001$; digging: $z = 4.07$, $P < 0.001$; hind-kick: $z = 3.94$, $P < 0.001$). We also found that, once underground, males stayed in the burrow 28% longer than females (Fig. 3E; LMM, underground: $t = 2.86$, $P = 0.008$), but the mean duration of a digging bout was not significantly different between sexes (LMM, digging: $t = 0.25$, $P = 0.804$). Overall, we found that males spent 74% more time underground and 42% more time digging than females (Fig. 3F; LMMs, underground: $t = 4.54$, $P < 0.001$; digging: $t = 2.97$, $P = 0.006$). Together, these findings are consistent with the observation that individual males dig longer burrows than individual females.

We next examined the relative contributions of individual mice during pair trials. In same-sex pairs, the division of labor was statistically indistinguishable from 50/50, with each mouse contributing $50.0 \pm 3.7\%$ of the total digging duration for the pair (Fig. S3D; t -test, $t = 0.00$, $P = 1$). By contrast, the division of labor was significantly skewed in opposite-sex pairs, with males contributing $65.8 \pm 3.7\%$ of the total digging duration and females contributing $34.2 \pm 3.7\%$ (LMM, $F = 35.29$, $P < 0.001$). Because males spend more time digging than




Figure 3. Novel behavioral assay reveals sex and pair-type differences in burrowing behavior in *Peromyscus polionotus*. (A) Schematic of the narrow, transparent enclosures (left) and still image of two mice digging in tandem (right). (B) Experimental design. Baseline individual burrowing output was first quantified in virgin females and males (left). Mice were then tested, in random order, with a same-sex (middle) and opposite-sex (right) partner. (C) Total burrow length produced in three social contexts: individual, same-sex, and opposite-sex. Each line represents the mean of two trials per mouse, per context. (D) Number of underground, digging, and hind-kicking bouts. (E) Mean duration of underground and digging bouts. (F) Total time spent underground and total time spent digging. (D–F) Data reflect the mean of eight observations per individual, pooled across same-sex and opposite-sex trials. (G) Number of underground, digging, and hind-kicking bouts, across same-sex and opposite-sex trials. (H) Mean duration of underground and digging bouts, across trial-types. (I) Total time spent underground and total time spent digging, across trial-types. (G–I) Data reflect the mean of four observations per individual, per trial-type. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Error bars represent SEM.

females, we might expect, under a simple additive model, that male-male pairs would spend the most time digging and produce the longest burrows. However, we found that opposite-sex pairs spent just as much time digging (Fig. S3E; Tukey contrasts, MM

vs. FM: $t = 0.93$, $P = 0.625$) and produced burrows that were just as long as male-male pairs (Fig. S3C; Tukey contrasts, MM vs. FM: $t = 0.01$, $P = 1$). These findings suggest that digging behavior varies with social context in one or both sexes.

EFFECT OF SOCIAL CONTEXT ON INDIVIDUAL-LEVEL BURROWING BEHAVIOR

To determine how female and male *P. polionotus* might modulate their behavior according to social context, we compared how individuals behave across same-sex and opposite-sex trials. We used GLMMs with a Poisson link function to analyze “number of bouts” and LMMs to analyze “mean bout duration” and “total duration” per 10-minute observation period and tested for an effect of sex, social context, and sex by social context interaction (Table S1). First, we found no change in the number of underground bouts between same-sex and opposite-sex trials (Fig. 3G; LMM, $z = 1.61$, $P = 0.107$). However, females had 21% fewer digging bouts (planned contrasts, $z = 3.61$, $P < 0.001$) and performed 16% fewer hind-kicks ($z = 8.23$, $P < 0.001$) in opposite-sex trials, whereas males performed 53% more hind-kicks in opposite-sex trials ($z = 12.46$, $P < 0.001$). Second, for females in opposite-sex trials, we found a 36% increase in the mean duration of an underground bout (Fig. 3H; planned contrasts, $t = 3.96$, $P < 0.001$), but no change in the mean duration of a digging bout between same-sex and opposite-sex trials, for either sex (LMM, $t = 0.25$, $P = 0.804$). Third, females spent slightly (6%) more time underground in opposite-sex trials (Fig. 3I; planned contrasts, $t = 2.37$, $P = 0.019$). In addition, although females spent slightly less time digging in opposite-sex trials (same: 1.6 ± 0.3 min, opp: 1.2 ± 0.1) and males spent slightly more (same: 1.8 ± 0.2 min, opp: 2.2 ± 0.2), neither trend was significant (Fig. 3I). Together, these results suggest that, although some behaviors (e.g., number of bouts, mean bout duration) may differ slightly by social context, there was no strong difference in total digging duration by individuals in same-sex versus opposite-sex trials. Thus, mechanisms other than increased individual effort (i.e., total digging duration) are important for determining pair burrow length.

DIFFERENCES IN SOCIAL INTERACTION BETWEEN PAIR-TYPES

We next explored the frequency of affiliative and agonistic interactions that might indicate differences in social cohesion between same- and opposite-sex pairs. We calculated the cumulative number of affiliative interactions for a given pair across all four observation periods. Opposite-sex pairs engaged in 31% more affiliative behavior (e.g., allogrooming, huddling) than same-sex pairs (Fig. 4A; GLM, $z = 2.69$, $P = 0.007$). Next, because agonistic interactions were rarer than affiliative interactions, we calculated the fraction of trials in which any agonistic behavior was observed between mice. We recorded agonistic interactions (e.g., boxing, biting, fleeing) in 44% of male-male trials, but only 28% of opposite-sex trials, and no agonistic interactions were observed in any female-female trial (Fig. 4B; Fisher’s exact tests, FF vs. MM: $P = 0.007$; FF vs. FM: $P = 0.041$; MM vs. FM:




Figure 4. Opposite-sex *Peromyscus polionotus* pairs show more affiliative and fewer agonistic interactions than same-sex pairs. (A) Total number of affiliative interactions observed per pair. Each data point represents the sum of four observation periods across two trials. (B) Fraction of trials in which any agonistic behavior was observed. * $P < 0.05$, ** $P < 0.01$. Error bars represent SEM.

$P = 0.339$). These results raise the possibility that differences in social cohesion between pair-types may reflect differences in the propensity to burrow cooperatively.

DIFFERENCES IN DIGGING EFFICIENCY BETWEEN PAIR-TYPES

To understand how opposite-sex pairs produce longer burrows despite no significant increase in total digging duration, we next tested for differences in digging efficiency between pair-types. To determine digging efficiency (i.e., burrow extension rate), we compared the change in burrow length over the course of each 10-minute observation period to the total digging duration observed for both mice during that same period (see *Methods* in the Supporting Information). We distinguished between independent digging (i.e., mice working independently, either temporally or spatially) and simultaneous digging (i.e., mice working together, in the same burrow at the same time) (Video S1). To control for the number of mice, independent and simultaneous digging were both expressed in person-hours or, more accurately, mouse-minutes (i.e., time spent digging multiplied by the number of mice digging). To test for an effect of total digging duration on change in burrow length, we used an LMM with independent digging and simultaneous digging as fixed effects and pair ID, observer ID, trial number, and observation period as random effects. As expected, the durations of both independent digging and simultaneous digging were significant predictors of change in burrow length (LMM, independent: $t = 3.18$, $P = 0.002$;




Figure 5. Opposite-sex pairs are more likely to engage in more efficient simultaneous digging. (A, B) Partial regression plots. (A) Estimated relationship between independent digging duration (i.e., one mouse working at a time) and change in burrow length, controlling for simultaneous digging duration. (B) Estimated relationship between simultaneous digging duration (i.e., both mice working together) and change in burrow length, controlling for independent digging duration. (C) Effect estimates for independent and simultaneous digging duration (in mouse-minutes) on change in burrow length. (D) Probability of observing simultaneous digging for a range of simulated focal mouse digging durations, given a fixed partner mouse digging duration. (E) Probability of observing simultaneous digging for same-sex and opposite-sex pairs. * $P < 0.05$. Shaded areas and error bars represent 95% confidence intervals.

simultaneous: $t = 5.20$, $P < 0.001$). Next, to compare the efficiency of these two digging modes, we calculated partial correlation coefficients for both independent and simultaneous digging (Fig. 5A; independent digging: $r = 0.255$, $P < 0.001$, Fig. 5B; simultaneous digging: $r = 0.356$, $P < 0.001$). We found that an additional mouse-minute of independent digging resulted in an additional 0.46 ± 0.13 cm of burrow length (Fig. 5C). By contrast, an additional mouse-minute of simultaneous digging resulted in an additional 1.09 ± 0.21 cm of burrow length (Fig. 5C), indicating that simultaneous digging is, on average, 135% more efficient than independent digging, even when controlling for the number of mice digging.

We then asked if the probability of engaging in more efficient simultaneous digging differed between pair-types. Although both same-sex and opposite-sex pairs engaged in both digging modes, opposite-sex pairs were more likely to engage in simultaneous digging, even when controlling for the total digging duration for both mice (Fig. 5D; GLM, $z = 2.24$, $P = 0.025$). The probability of observing simultaneous digging was significantly higher for opposite-sex than for same-sex pairs: $54 \pm 5\%$ for opposite-sex pairs, but only $32 \pm 8\%$ for female-female pairs and $39 \pm 7\%$ for male-male pairs (Fig. 5E). Together, our results suggest that longer opposite-sex burrows in *P. polionotus* can be explained predominately by an increase in burrowing

efficiency—likely mediated by enhanced social cohesion—rather than by an increase in individual burrowing effort.

Discussion

Here, we show, under controlled laboratory conditions, that social context differentially affects the expression of innate burrowing behavior in three species of *Peromyscus* mice. In two species—*P. leucopus* and *P. maniculatus*—we found no change in burrow length according to social context: the burrows built by individual mice were similar to those built when two mice were assayed together. By contrast, in *P. polionotus*, we found that pairs of mice jointly constructed burrows that were nearly twice as long as those dug by individuals. We further demonstrate, in *P. polionotus*, that opposite-sex pairs dig longer burrows than same-sex pairs, driven largely by an increase in digging efficiency (likely due to increased social cohesion) among opposite-sex pairs. Together, we find evidence of cooperative burrowing behavior in *P. polionotus*, but not its closest relatives.

Collective building behavior has evolved in diverse taxa—from social spiders (Avilés 1997) and eusocial insects (Theraulaz et al. 1998) to communally nesting birds (Cockburn 1998) and cooperatively burrowing mole-rats (Jarvis et al. 1994). Our observation, that burrows built by *P. polionotus* pairs are nearly twice as long as those built by individuals, is consistent with findings in other species that build collective structures. For example, in most ants and termites, nest volume correlates strongly with colony size (Perna and Theraulaz 2017). This adaptive scaling is mediated by a negative-feedback process in which insects adjust their digging rate according to traffic flow or the density of workers in the nest (Rasse and Deneubourg 2001). Similarly, bird species in which both sexes contribute to nest building have larger nests than those in which only the female builds the nest (Soler et al. 1998). Thus, in many species, structures built by pairs or groups are larger than structures built by individuals—a trend that we observed in *P. polionotus*, but not *P. leucopus* or *P. maniculatus*. Perhaps even more interestingly, in *P. polionotus*, burrow length differed according to not only the number of diggers (i.e., individuals vs. pairs) but also the composition of the digging pair (i.e., same-sex vs. opposite-sex).

Within *P. polionotus*, both same- and opposite-sex pairs cooperatively constructed burrows, but those dug by opposite-sex pairs were considerably longer. By designing a novel behavioral assay, we could explicitly quantify individual burrowing behavior across different social contexts to determine whether the observed difference in burrow length between same-sex and opposite-sex pairs was due to changes in digging duration, digging efficiency, or both. Surprisingly, neither males nor females spent significantly more time digging with an opposite-sex partner, but instead, longer opposite-sex burrows arose largely due to

increased digging efficiency (i.e., an increased rate of burrow extension). Specifically, opposite-sex pairs were more likely to dig simultaneously in the same burrow, perhaps in part because they were more socially cohesive and tolerant of a digging partner. Simultaneous digging appeared to be a more efficient mode of burrow extension than two mice digging independently. The simultaneous digging behavior we observed in *P. polionotus* pairs is reminiscent of “chain digging” observed in communally burrowing mammals, such as eusocial mole-rats (Lovegrove 1989) and group-living degus (Ebensperger and Bozinovic 2000), and suggests that even distantly related rodent species may have converged upon a common strategy for efficiently excavating shared living space.

One important unanswered question is whether the longer burrows dug by opposite-sex pairs confer fitness benefits to the pair and/or their offspring. For example, a longer burrow may provide additional protection from predators or buffer against temperature and humidity fluctuations, which are particularly harmful to pups (Berry and Bronson 1992; Bedford et al. 2021). Thus, reproductive pairs may engage in more simultaneous digging to efficiently construct longer burrows, which may, in turn, increase the odds of survival for the pair and/or their pups. Alternatively, longer opposite-sex burrows could arise simply as a by-product of increased social cohesion between opposite-sex partners, with little or no fitness benefits. Notably, the prospect of future reproduction is not strictly necessary for cooperative burrowing behavior in this species: even same-sex *P. polionotus* pairs constructed longer burrows than individuals, even if not as long as opposite-sex burrows. Indeed, in the wild, two adult females or two adult males are occasionally found sharing a single burrow (Rand and Host 1942; Blair 1951; Smith 1966), and survival benefits may also accrue to same-sex pairs that cooperatively construct long burrows. Thus, additional research quantifying the consequences of burrow length variation for both survival and reproduction in the wild is needed to fully untangle the selective pressures (if any) driving the evolution of cooperative burrowing in *Peromyscus*.

Our study revealed that only the monogamous *P. polionotus* demonstrated cooperative burrowing behavior. Although an association between monogamy and cooperation has been previously noted (Dillard and Westneat 2016; Hahn et al. 2021), in this study, these species also differ ecologically, and therefore we cannot rule out the contributions of such factors to interspecific variation in cooperative burrowing. For example, both *P. leucopus* and *P. maniculatus* occupy structured habitats, such as forests and grasslands, and nest in a variety of locations, including rock crevices, brush piles, fallen logs, and tree cavities (Nicholson 1941; Madison et al. 1984; Sharpe and Millar 1990). This diversity of alternative refuges suggests that these mice may only infrequently excavate a burrow *de novo*. In addition,

radiotelemetry (Mineau and Madison 1977; Wolff and Hurlbutt 1982; Madison et al. 1984) and nest-box studies (Nicholson 1941; Wolff and Durr 1986) indicate a general pattern of solitary nesting. Thus, *P. leucopus* and *P. maniculatus* may be facultative, solitary burrowers with few opportunities for cooperative burrowing in the wild. By contrast, *P. polionotus* occupies open habitats, such as fallow fields and sand dunes, and nests almost exclusively in burrows of its own construction (Blair 1951; Weber et al. 2013; Hu and Hoekstra 2017). Moreover, *P. polionotus* often nests in opposite-sex pairs (Rand and Host 1942; Blair 1951; Smith 1966) and is likely an obligate burrower with ample opportunity for cooperative burrowing in the wild. Thus, in addition to social organization, ecological factors such as habitat type and the availability of alternative nest sites could play a role in the evolution of cooperative burrowing in *Peromyscus*. Further studies examining ecological factors known to modulate behavioral flexibility (e.g., habitat complexity [Leal and Powell 2012], predation risk [Ghalambor et al. 2013], etc.) are necessary to test this hypothesis.

Finally, our results suggest that different *Peromyscus* species may follow different decision rules that govern their burrowing behavior. *Peromyscus maniculatus* and *P. leucopus*, for example, may employ an extrinsic, goal-oriented strategy in which mice dig until sufficient habitable space has been excavated, resulting in similarly sized burrows regardless of social context (i.e., digging alone or in pairs). By contrast, *P. polionotus* may employ an intrinsic, effort-oriented strategy in which mice dig until a certain amount of individual effort has been expended, resulting in significantly longer burrows when mice are paired. Our results offer new hypotheses for how different decision rules for building behavior might arise among closely related species.

Overall, our study demonstrates that social context can affect the expression of innate behavior in different ways, even among closely related species. We also show that variation in burrow length can arise in surprising ways: longer opposite-sex burrows in *P. polionotus* arose due to increased digging efficiency rather than increased individual effort. Together, these findings underscore the importance of combining direct observations of building behavior with measurements of the resulting structures to unravel mechanisms driving behavioral flexibility.

ACKNOWLEDGMENTS

We thank N. Shen Molesky for help with behavioral assays, Z. Ali for helpful discussions, and E. Hager, O. Harringmeyer, C. Hu, and C. Lewarch for comments on the manuscript. This work was supported by a James Mills Pierce Fellowship, Robert A. Chapman Memorial Scholarship, and a Natural Science and Engineering Research Council of Canada Postgraduate Scholarship to NLB; a Robert A. Chapman Memorial Scholarship and a Harvard Mind Brain Behavior Graduate Student Award to JNW; a Harvard Mind Brain Behavior Graduate Student Award to WT; an HHMI International Student Research Fellowship, Herchel

Smith Graduate Fellowship, and a Joan Brockman Williamson Graduate Fellowship to FB; a Harvard College Program for Research in Science and Engineering Fellowship to AK; and a Harvard College Program for Research in Science and Engineering Fellowship and Herchel Smith Undergraduate Fellowship to RAG. HEH is an Investigator of the Howard Hughes Medical Institute.

AUTHOR CONTRIBUTIONS

NLB, JNW, WT, FB, and HEH designed the experiments. NLB, JNW, WT, FB, AK, and RAG performed the experiments. NLB analyzed the data. NLB and HEH wrote the manuscript with input from all authors.

CONFLICT OF INTERESTS

The authors declare no conflict of interest.

REFERENCES

- Amici, F., Aureli, F. & Call, J. (2008). Fission-fusion dynamics, behavioral flexibility, and inhibitory control in primates. *Current Biology*, 18, 1415–1419.
- Avilés, L. (1997). Causes and consequences of cooperation and permanent-sociality in spiders. pp. 476–498 in J. C. Choe & B. J. Crespi, eds. *The evolution of social behaviour in insects and arachnids*. Cambridge Univ. Press, Cambridge U.K.
- Bedford, N. L., Gable, J. T., Hu, C. K., Wooldridge, T. B., Sokolov, N. A., Lassance, J.-M., et al. (2021). Automated tracking reveals the social network of beach mice and their burrows. *bioRxiv*, <https://doi.org/10.1101/2021.08.07.455531>
- Bendesky, A., Kwon, Y.-M., Lassance, J.-M., Lewarch, C. L., Yao, S., Peterson, B. K., et al. (2017). The genetic basis of parental care evolution in monogamous mice. *Nature*, 544, 434–439.
- Berry, R. J. & Bronson, F. H. (1992). Life history and bioeconomy of the house mouse. *Biological Reviews*, 67, 519–550.
- Blair, W. F. (1951). Population structure, social behavior, and environmental relations in a natural population of the beach mouse (*Peromyscus polionotus leucocephalus*). *Contributions from the Laboratory of Vertebrate Biology*, 48, 1–47.
- Bond, A. B., Kamil, A. C. & Balda, R. P. (2007). Serial reversal learning and the evolution of behavioral flexibility in three species of North American corvids (*Gymnorhinus cyanocephalus*, *Nucifraga columbiana*, *Aphelocoma californica*). *Journal of Comparative Psychology*, 121, 372–379.
- Cockburn, A. (1998). Evolution of helping behavior in cooperatively breeding birds. *Annual Review of Ecology and Systematics*, 29, 141–177.
- Dawson, W. D., Lake, C. E. & Schumpert, S. S. (1988). Inheritance of burrow building in *Peromyscus*. *Behavior Genetics*, 18, 371–382.
- Dillard, J. R. & Westneat, D. F. (2016). Disentangling the correlated evolution of monogamy and cooperation. *Trends in Ecology & Evolution*, 31, 503–513.
- Ebensperger, L. A. & Bozinovic, F. (2000). Communal burrowing in the hystriognath rodent, *Octodon degus*: a benefit of sociality? *Behavioral Ecology and Sociobiology*, 47, 365–369.
- Foltz, D. W. (1981). Genetic evidence for long-term monogamy in a small rodent, *Peromyscus polionotus*. *The American Naturalist*, 117, 665–675.
- Ghalambor, C. K., Peluc, S. I. & Martin, T. E. (2013). Plasticity of parental care under the risk of predation: how much should parents reduce care? *Biology Letters*, 9, 20130154.
- Gonzalez-Mariscal, G., McNitt, J. I. & Lukefahr, S. D. (2007). Maternal care of rabbits in the lab and on the farm: endocrine regulation of behavior and productivity. *Hormones and Behavior*, 52, 86–91.

- Guayasamin, O. L., Couzin, I. D. & Miller, N. Y. (2017). Behavioural plasticity across social contexts is regulated by the directionality of inter-individual differences. *Behavioural Processes*, 141, 196–204.
- Hahn, L. G., Hooper, R., McIvor, G. E. & Thornton, A. (2021). Cooperative nest building in wild jackdaw pairs. *Animal Behaviour*, 178, 149–163.
- Hoffmann, A. A. (1990). The influence of age and experience with conspecifics on territorial behavior in *Drosophila melanogaster*. *Journal of Insect Behavior*, 3, 1–12.
- Hu, C. K. & Hoekstra, H. E. (2017). *Peromyscus* burrowing: a model system for behavioral evolution. *Seminars in Cell & Developmental Biology*, 61, 107–114.
- Jarvis, J. U., O’Riain, M. J., Bennett, N. C. & Sherman, P. W. (1994). Mammalian eusociality: a family affair. *Trends in Ecology & Evolution*, 9, 47–51.
- Kappeler, P. M., Barrett, L., Blumstein, D. T. & Clutton-Brock, T. H. (2013). Constraints and flexibility in mammalian social behaviour: introduction and synthesis. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368, 20120337.
- Leal, M. & Powell, B. J. (2012). Behavioural flexibility and problem-solving in a tropical lizard. *Biology Letters*, 8, 28–30.
- Lovegrove, B. (1989). The cost of burrowing by the social mole rats (Bathyergidae) *Cryptomys damarensis* and *Heterocephalus glaber*: the role of soil moisture. *Physiological Zoology*, 62, 449–469.
- Madison, D., Hill, J. & Gleason, P. (1984). Seasonality in the nesting behavior of *Peromyscus leucopus*. *American Midland Naturalist*, 112, 201–204.
- Mayr, E. (1974). Behavior programs and evolutionary strategies: natural selection sometimes favors a genetically “closed” behavior program, sometimes an “open” one. *American Scientist*, 62, 650–659.
- Metz, H. C., Bedford, N. L., Pan, Y. L. & Hoekstra, H. E. (2017). Evolution and genetics of precocious burrowing behavior in *Peromyscus* mice. *Current Biology*, 27, 3837–3845. e3833.
- Mineau, P. & Madison, D. (1977). Radio-tracking of *Peromyscus leucopus*. *Canadian Journal of Zoology*, 55, 465–468.
- Montiglio, P. O., Wey, T. W., Chang, A. T., Fogarty, S. & Sih, A. (2017). Correlational selection on personality and social plasticity: morphology and social context determine behavioural effects on mating success. *Journal of Animal Ecology*, 86, 213–226.
- Nelson, A. C., Cauceglia, J. W., Merkle, S. D., Youngson, N. A., Oler, A. J., Nelson, R. J. *et al.* (2013). Reintroducing domesticated wild mice to sociality induces adaptive transgenerational effects on MUP expression. *Proceedings of the National Academy of Sciences*, 110, 19848–19853.
- Nicholson, A. J. (1941). The homes and social habits of the wood-mouse (*Peromyscus leucopus noveboracensis*) in southern Michigan. *American Midland Naturalist*, 25, 196–223.
- Perna, A. & Theraulaz, G. (2017). When social behaviour is moulded in clay: on growth and form of social insect nests. *Journal of Experimental Biology*, 220, 83–91.
- Ramdy, P., Lichocki, P., Cruchet, S., Frisch, L., Tse, W., Floreano, D. *et al.* (2015). Mechanosensory interactions drive collective behaviour in *Drosophila*. *Nature*, 519, 233–236.
- Rand, A. & Host, P. (1942). Mammal notes from Highlands County, Florida; results of the Archbold Expeditions. *Bulletin of the American Museum of Natural History*, 81, 1–21.
- Rasse, P. & Deneubourg, J.-L. (2001). Dynamics of nest excavation and nest size regulation of *Lasius niger* (Hymenoptera: Formicidae). *Journal of Insect Behavior*, 14, 433–449.
- Ribble, D. O. & Millar, J. S. (1996). The mating system of northern populations of *Peromyscus maniculatus* as revealed by radiotelemetry and DNA fingerprinting. *Ecoscience*, 3, 423–428.
- Royle, N. J., Russell, A. F. & Wilson, A. J. (2014). The evolution of flexible parenting. *Science*, 345, 776–781.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T. *et al.* (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, 9, 676–682.
- Sharpe, R. (1975). The influence of the sex of litter-mates on subsequent maternal behaviour in *Rattus norvegicus*. *Animal Behaviour*, 23, 551–559.
- Sharpe, S. T. & Millar, J. (1990). Relocation of nest sites by female deer mice, *Peromyscus maniculatus borealis*. *Canadian Journal of Zoology*, 68, 2364–2367.
- Smith, M. H. (1966). The evolutionary significance of certain behavioral, physiological, and morphological adaptations of the old-field mouse, *Peromyscus polionotus*. Ph.D. diss., University of Florida, Gainesville, FL.
- Soler, J. J., Moller, A. P. & Soler, M. (1998). Nest building, sexual selection and parental investment. *Evolutionary Ecology*, 12, 427–441.
- Stagkourakis, S., Spigolon, G., Liu, G. & Anderson, D. J. (2020). Experience-dependent plasticity in an innate social behavior is mediated by hypothalamic LTP. *Proceedings of the National Academy of Sciences*, 117, 25789–25799.
- Taborsky, B., Arnold, C., Junker, J. & Tschopp, A. (2012). The early social environment affects social competence in a cooperative breeder. *Animal Behaviour*, 83, 1067–1074.
- Theraulaz, G., Bonabeau, E. & Deneubourg, J. L. (1998). The origin of nest complexity in social insects. *Complexity*, 3, 15–25.
- Watanabe, K., Chiu, H., Pfeiffer, B. D., Wong, A. M., Hooper, E. D., Rubin, G. M. *et al.* (2017). A circuit node that integrates convergent input from neuromodulatory and social behavior-promoting neurons to control aggression in *Drosophila*. *Neuron*, 95, 1112–1128.
- Weber, J. N. & Hoekstra, H. E. (2009). The evolution of burrowing behaviour in deer mice (genus *Peromyscus*). *Animal Behaviour*, 77, 603–609.
- Weber, J. N., Peterson, B. K. & Hoekstra, H. E. (2013). Discrete genetic modules are responsible for complex burrow evolution in *Peromyscus* mice. *Nature*, 493, 402–405.
- Wolff, J. O. & Durr, D. S. (1986). Winter nesting behavior of *Peromyscus leucopus* and *Peromyscus maniculatus*. *Journal of Mammalogy*, 67, 409–412.
- Wolff, J. O. & Hurlbutt, B. (1982). Day refuges of *Peromyscus leucopus* and *Peromyscus maniculatus*. *Journal of Mammalogy*, 63, 666–668.
- Xia, X. & Millar, J. S. (1988). Paternal behavior by *Peromyscus leucopus* in enclosures. *Canadian Journal of Zoology*, 66, 1184–1187.
- Xia, X. & Millar, J. S. (1991). Genetic evidence of promiscuity in *Peromyscus leucopus*. *Behavioral Ecology and Sociobiology*, 28, 171–178.
- R Core Team (2021). R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. URL <https://www.R-project.org/>

Associate Editor: Dr. C. Moreau

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Supplementary Figures

Supplementary Methods 2

Table S1

Video S1: Examples of independent and simultaneous digging in pairs of *P. polionotus* mice

EVOLUTION LETTERS 2022

11

Supplementary Methods

Animal husbandry

We performed experiments with three *Peromyscus* species: *P. polionotus subgriseus* (PO), *P. maniculatus bairdii* (BW), and *P. leucopus* (LL) obtained from the *Peromyscus* Genetic Stock Center at the University of South Carolina but maintained as outbred colonies at Harvard University. We housed animals in ventilated cages measuring 7.75 x 12 x 6.5" (Allentown Inc., Allentown, NJ), which were furnished with 1/4" Bed-o'Cobs bedding (The Andersons, Maumee, OH), a 2" square cotton nestlet (Ancare, Bellmore, NY), and a red polycarbonate hut (Bio-Serv, Flemington, NJ). Mice were provided food and water *ad libitum*. Upon weaning at 23 days of age, mice were co-housed in same-sex groups of up to five conspecifics and fed irradiated Prolab Isopro RMH 3000 (LabDiet, St. Louis, MO). We fed all paired adults irradiated breeder chow: PicoLab Mouse Diet 20 (LabDiet, St. Louis, MO). We maintained mice on a 16h light:8h dark cycle at 22°C. The Institutional Animal Care and Use Committee at Harvard University approved all protocols.

Behavioral assays

We measured innate burrowing behavior in two complementary assays under controlled laboratory conditions, allowing us to minimize the effects of other environmental variables (which could also contribute to, for example, interspecific variation in behavioral flexibility).

Large enclosures. We measured burrow architecture as described previously (Weber & Hoekstra 2009; Weber *et al.* 2013; Metz *et al.* 2017). Briefly, at the start of the dark cycle, we introduced individuals or pairs of mice into large PVC boxes (1.2 x 1.5 x 1.1 m) filled with approximately 700 kg hydrated, hard-packed premium play sand (Quickrete, Atlanta, GA). In the enclosures, mice were provided food and water *ad libitum* along with a cotton nestlet. We removed mice from the enclosures after one (pregnancy trials) or two (standard trials) overnight periods. Then, we made casts of the resultant burrows using polyurethane filling foam (Hilti, Schaan, Liechtenstein). Next, a researcher blind to trial identity hand-measured the burrow casts. For each trial, the number of burrows and total length of each burrow was recorded. We also categorized burrow shape as either "simple" or "complex" as described previously (Weber & Hoekstra 2009): *P. leucopus* and *P. maniculatus* burrows are considered "simple" with a short entrance tunnel and terminal nest chamber, whereas *P. polionotus* burrows are considered "complex" with a long entrance tunnel, nest chamber, and escape tunnel. After each trial, we removed all food, feces, nesting material, and disturbed substrate from the enclosure, and then rinsed the enclosure walls with water and turned over the sand to minimize residual odors.

Narrow, transparent enclosures. We also assayed burrowing behavior of mice directly using an acrylic chamber (5 x 90 x 60 cm) with a transparent Plexiglass face (see Fig. 3A). Using a pre-cut mold, we sculpted hydrated sand into two symmetrical 45° hills and excavated an 8 cm tunnel from one randomly selected hill to encourage burrowing in a consistent location. We outfitted the apparatus with an infrared illuminator frame that enabled video recording in the dark. We then introduced individuals or pairs of mice into an enclosure at the start of the dark cycle and removed animals the next morning, recording 8 hours of continuous video during the dark cycle. Mice were provided food and water *ad libitum*. Photographs of the resultant burrows were taken at the end of the trial. Details of photo and video analyses

are provided below. Following each trial, we removed all food, feces, and disturbed substrate and rinsed the enclosure to minimize residual odors.

Experimental design

Effect of social context on burrow length. In the large enclosures, we first measured the burrows of mice from three *Peromyscus* species in individual and pair trials. Sample sizes are indicated in Figure 1. To control for past experience, we randomized the order in which mice were tested in individual, same-sex, or opposite-sex trials. Based on mouse availability, we tested 25 mice once, 43 twice, and 25 more than twice. For pair trials, we tested 34 pairs once, 48 twice, and 2 more than twice. On average, each individual, or each unique pair, was tested 1.6 times (range: 1–7 trials). For all trials, mice were released into the enclosures at the start of the dark cycle and retrieved after two overnight periods (~32 hours).

Effect of reproductive state on female burrow length. To test whether pairing with a male or pregnancy alters burrowing performance, we compared the burrows dug by females before and after being co-housed with a male. First, we tested the burrowing output of virgin females ($n = 42$ *P. polionotus*, $n = 36$ *P. maniculatus*) in the large enclosures. Each female was assayed twice with 2 days rest between trials. We then transferred each female to a new cage with a conspecific male. Approximately half of all paired females subsequently became pregnant (21/42 *P. polionotus*, 19/36 *P. maniculatus*). Co-housed females were then tested again twice, as individuals, with 2 days rest between trials. Females were returned to their home cage after each trial and monitored daily for parturition for up to 23 days. To minimize stress to pregnant females, all mice in this experiment were tested for only one overnight period (~8 hours) in the large enclosures.

Effect of social context on individual-level burrowing behavior. To test whether *P. polionotus* mice modulate their behavior according to social context, we used a repeated-measures design to test mice in individual, same-sex, and opposite-sex trials in the narrow, transparent enclosures. To distinguish individuals during pair trials, we shaved a patch of hair from both flanks of one randomly selected member of the pair. Shaving was completed at least a week before the first pairing and was visible over the course of the experiment. We excluded behavioral data from one same-sex pair (FF) in which the markings of the two females were indistinguishable and mouse ID could not be confidently assigned. To quantify baseline burrowing output, we first assayed each individual twice ($n = 16$ virgin females, $n = 16$ virgin males). Next, we transferred each individual to a new cage with an unrelated, unfamiliar partner of the same or opposite sex. Pairs were given 2 nights to acclimate before being tested together in the transparent enclosures. We recorded 8 hours of video for each overnight trial, and pairs were then returned to their home cage for 3 nights rest before their second trial. Mice were then re-partnered and the process was repeated. To control for previous experience, we randomly assigned the order in which mice burrowed with a same- or opposite-sex partner. Females were monitored daily for parturition for up to 23 days after the end of the experiment; no females were pregnant at any point during the experiment.

Behavioral analyses

Photo analysis. To measure the length of burrows dug by individuals and pairs of mice in the narrow, transparent enclosures, we took photos of the enclosures at the end of the 8h trial. Each image was measured separately by two researchers, both blind to trial identity, using Fiji image processing software (Schindelin *et al.* 2012). Burrow length measurements were highly correlated between researchers

(Pearson correlation, $r = 0.93$, $P < 0.001$); therefore, we used the mean of these two measurements for all statistical analyses. We also took video stills at the beginning and end of each 10-minute observation period (see *Video analysis*). From these stills, we measured burrow length as described above and subtracted the initial burrow length from the final burrow length to calculate change in burrow length over the 10-minute observation period.

Video analysis. In 75% of trials, the mice had fully completed at least one burrow 3 hours after the start of the dark cycle. Thus, to target the active burrow construction phase, we selected 2 10-minute observation periods, spaced one hour apart: 0:30–0:40 and 1:30–1:40 hours. In 5/62 videos, we did not observe any digging during either of the two observation periods. For these videos, we shifted our two observation periods, still spaced one hour apart, to slightly later in the first 3-hour time window. We accounted for this shift in our statistical analyses by modeling observation period as a random effect (see *Statistics*).

We quantified the behavior of each mouse in a pair using The Observer XT Version 12.0 (Noldus, Leesburg, VA). Each 10-minute observation period ($n = 124$) was randomly assigned to a researcher blind to trial identity for behavioral scoring according to the following scheme: for each individual, we scored all burrow entries or exits (to calculate time spent underground), as well as extending (i.e., forelimb digging at the growing end of the burrow), widening (i.e., forelimb digging at any other location in the burrow), and hind-kicks (i.e., vigorous, coordinated hindlimb movements that expel loosened sand from the burrow). Sand would occasionally stick to the Plexiglass face of the enclosure, temporarily obscuring the mouse in the burrow; we noted all instances where the mouse was underground, but obscured, thereby precluding our ability to score other behaviors, such as digging or social behavior. We found no effect of social context on how often mouse behavior was obscured (LMM, $t = 1.57$, $P = 0.124$). Time spent underground, time spent obscured, burrow extending, and burrow widening were all scored as state events, whereas hind-kicks were scored as point events. Point events were analyzed as counts only, whereas state events were analyzed as both counts and durations. All events were recorded in Observer to the nearest hundredth of a second. For each pair, we also scored affiliative and agonistic behaviors, which could occur either inside or outside the burrow. We defined affiliative behaviors as beneficial social interactions (i.e., allogrooming, huddling) and agonistic behaviors as aggressive interactions (i.e., boxing, parrying, biting) or submissive behaviors (i.e., freezing, fleeing). Affiliative and agonistic behaviors were scored as state events. Last, to determine how often two mice dug in the same burrow at the same time, we aligned the digging ethograms of the two mice and calculated all instances of overlap. Thus, for each observation period, we calculated the total time spent digging alone and the total time spent digging together.

Altogether, each mouse ($n = 32$) received 80 minutes of direct observation across the entire study: 10 minutes per observation period, 2 observation periods per trial, 2 trials per social context, and 2 social contexts (i.e., same-sex and opposite-sex trials).

Statistics

All statistical tests were performed in R Version 4.1.1 (R Core Team, 2021). We used the *lmer* and *glmer* functions (lme4 package) to run linear mixed-effects models (LMMs) and generalized linear mixed-effects models (GLMMs), respectively. We used the *anova* and *summary* functions (lmerTest package) to determine degrees of freedom, test statistics, and p-values for fixed effect terms. We used the *emmeans*

and *pairs* functions (emmeans package) to determine degrees of freedom, test statistics, and p-values for planned contrasts.

Effect of social context on burrow length. Using LMMs, we tested for an effect of species, social context (i.e., individual vs. pair trial), and species by social context interaction on “number of burrows” and “maximum burrow length” dug per trial. To improve normality, we performed a square-root transform on maximum burrow length. We included mouse sex as a fixed effect and mouse ID as a random effect. We calculated the pair: individual burrow length ratio by dividing the length of the average burrow dug by pairs of mice by the length of the average burrow dug by individual mice, excluding 5 *P. leucopus* individuals and 1 *P. leucopus* pair that dug no burrows. Using a Taylor series expansion, we then approximated the variance of each ratio. In *P. polionotus*, we used an LMM to test for an effect of trial-type on maximum burrow length. We modeled mouse ID as a random effect. We then used planned contrasts to test for differences in maximum burrow length between individual females and males, and between same-sex and opposite-sex pairs. Next, we calculated the observed: expected burrow length ratio for each unique pair of mice by dividing the average burrow length dug by a pair of mice by the sum of the average burrow lengths dug by individuals comprising that pair. Using an LMM, we then tested for an effect of pair-type (i.e., same-sex vs. opposite-sex pairs), modeling mouse ID and partner ID as random effects. We tested whether the observed: expected burrow length ratios for FF, MM, and FM pairs were significantly different from 1 using one sample *t*-tests with $\mu = 1.0$.

Effect of reproductive state on female burrow length. To first test for an effect of virgin burrow length on the probability of becoming pregnant, we used a GLMM with a binomial link function, modeling mouse ID as a random effect. We then used an LMM to test for an effect of cohabitation with a male, pregnancy, and cohabitation by pregnancy interaction on maximum burrow length, modeling mouse ID as a random effect in both *P. polionotus* and *P. maniculatus*. To improve normality, we log-transformed maximum burrow length. In *P. maniculatus*, we then used planned contrasts to test for an effect of cohabitation with a male on maximum burrow length in mice that became pregnant versus those that did not.

Sex differences in burrow length. Using LMMs, we tested for an effect social context (i.e., individual vs. pair trial) on “number of burrows” and “maximum burrow length” dug per trial in the narrow, transparent enclosures (see *Photo analysis*). We included mouse sex as a fixed effect and mouse ID as a random effect. We also used an LMM with mouse ID as a random effect to test for an effect of trial-type on maximum burrow length. We followed up with planned contrasts to test for differences between individual females and males, and between same-sex and opposite-sex pairs. Next, we tested for an effect of trial-type (i.e., individual, same-sex, or opposite-sex), sex, and trial-type by sex interaction on total burrow length. We used an LMM with mouse ID as a random effect and followed up with planned contrasts to test whether females and males respond differently to changes in social context.

Sex differences in burrowing behavior. Because we found no effect of sex or pair-type on the ratio of burrow extending to burrow widening (LMM, sex: $F = 0.25$, $P = 0.617$; pair-type: $F = 0.40$, $P = 0.528$), we collapsed these two categories into one general “digging” designation. We then tested for sex differences in underground and digging behavior. We pooled data for individual females and males across same-sex and opposite-sex trials and controlled for social context by including pair-type as a fixed effect in all models. We used GLMMs with Poisson link functions to analyze the number of behavior bouts and LMMs to analyze mean bout duration and total duration. To improve normality, we log-transformed mean bout duration and total duration. All models included mouse sex and pair-type as fixed effects and mouse

ID, observer ID, trial number, and observation period as random effects. To calculate the division of labor in each observation period, we divided the total digging duration for each mouse by the total digging duration for the pair. For same-sex pairs, we tested whether the division of labor was significantly different from 0.5 using a one sample t -test with $\mu = 0.5$. We tested whether the division of labor was skewed in opposite-sex pairs using an LMM with sex as a fixed effect and mouse ID, observer ID, trial number, and observation period as random effects. Last, we tested for pair-type differences in total digging duration using an LMM with pair-type as a fixed effect and pair ID, observer ID, trial number, and observation period as random effects. We followed up with Tukey contrasts to test for differences among FF, MM, and FM pairs.

Effect of social context on individual-level burrowing behavior. To assess whether individual mice behave differently in same-sex and opposite-sex trials, we tested for an effect of sex, social context, and sex by social context interaction on underground and digging behavior. We used GLMMs with Poisson link functions to analyze number of bouts and LMMs to analyze mean bout duration and total duration. To improve normality, we log-transformed mean bout duration and total duration. All models included mouse ID, observer ID, trial number, and observation period as random effects (Table S1). We removed all non-significant interaction effects from our models and followed up with planned contrasts to explore significant interaction effects.

Differences in social interaction between pair-types. To quantify differences in social cohesion between pair-types, we calculated the cumulative number of affiliative interactions for a given pair of mice across all 4 observation periods. Using a GLM with a Poisson link function, we tested for an effect of pair-type. Agonistic interactions were more rare than affiliative interactions. We therefore calculated the fraction trials in which any agonistic behavior was observed for a given pair of mice. We used Fisher's exact tests to test for differences among same-sex female-female (FF) and male-male (MM) pairs as well as opposite-sex female-male (FM) pairs.

Differences in digging efficiency between pair-types. To determine digging efficiency, we first measured the change in burrow length over each 10-minute observation period using still images from the video taken at the beginning and end of each observation period (see *Photo analysis*). We then calculated the total duration of both independent digging (i.e., two mice working independently, either temporally or spatially) and simultaneous digging (i.e., two mice working together, in the same burrow at the same time) during the observation period (see *Video analysis*). We expressed the total duration of each digging mode in "mouse-minutes" (i.e., time spent digging multiplied by the number of mice digging).

We tested for an effect of total digging duration on change in burrow length using an LMM with independent digging and simultaneous digging as fixed effects and pair ID, observer ID, trial number, and observation period as random effects. We used partial correlation tests (*pcor.test* function, *ppcor* package) to determine the correlation between independent digging and change in burrow length, while controlling for simultaneous digging, as well as the correlation between simultaneous digging and change in burrow length, while controlling for independent digging. Because the probability of overlap may increase with digging duration, we calculated the predicted probability of observing simultaneous digging at each value of total digging duration for the focal mouse (which ranged from 0 to 6.83 min), while holding the total digging duration for the partner mouse constant at its mean (1.49 min). We then tested whether the probability of observing simultaneous digging at a given burrow during a given observation period

differed between pair-types. We used a GLM with a binomial link function, including pair-type and total digging duration for both mice as fixed effects.

Table S1A

Model summaries for the number of bouts of underground, digging, and hind-kicking behavior.

Response variable (model)	Fixed effects	Estimate	SE	<i>z</i>	<i>P</i>
Underground (GLMM)	Intercept	12.13	1.24	11.81	< 0.001
	Sex	18.46	1.37	4.01	< 0.001
	Context	11.46	1.28	-1.61	0.107
Digging (GLMM)	Intercept	11.82	1.39	7.55	< 0.001
	Sex	16.40	1.60	2.31	0.021
	Context	9.24	1.48	-3.61	< 0.001
	Sex*Context	16.04	1.50	3.83	< 0.001
Hind-kicking (GLMM)	Intercept	19.21	1.63	6.07	< 0.001
	Sex	24.93	2.01	1.23	0.220
	Context	12.53	1.71	-8.23	< 0.001
	Sex*Context	41.16	1.73	13.04	< 0.001

GLMMs were run with Poisson link functions. The reference sex is “female” and the reference social context is “same-sex”. All models included mouse ID, observer ID, trial number, and observation period as random effects. Statistically significant results are shown in bold. Corresponds to Figure 3G.

Table S1B

Model summaries for the mean bout duration of underground and digging behavior.

Response variable (model)	Fixed effects	Estimate	SE	<i>df</i>	<i>t</i>	<i>P</i>
Underground (LMM)	Intercept	5.79	1.18	11.89	10.54	< 0.001
	Sex	11.64	1.39	64.10	4.23	< 0.001
	Context	9.93	1.35	215.88	3.99	< 0.001
	Sex*Context	3.25	1.42	210.00	-3.11	0.002
Digging (LMM)	Intercept	6.71	1.23	2.50	9.04	0.006
	Sex	7.11	1.35	32.53	0.66	0.516
	Context	6.83	1.32	193.37	0.25	0.804

LMMs were performed on log-transformed response variables. Back-transformed estimates and standard errors (SE) are reported. The reference sex is “female” and the reference social context is “same-sex”. All models included mouse ID, observer ID, trial number, and observation period as random effects. Statistically significant results are shown in bold. Corresponds to Figure 3H.

Table S1C

Model summaries for the total duration of underground and digging behavior.

Response variable (model)	Fixed effects	Estimate	SE	<i>df</i>	<i>t</i>	<i>P</i>
Underground (LMM)	Intercept	49.83	1.31	10.12	14.33	< 0.001
	Sex	173.52	1.70	69.75	4.86	< 0.001
	Context	85.52	1.65	209.88	2.39	0.018
	Sex*Context	26.72	1.79	208.09	-2.01	0.046
Digging (LMM)	Intercept	65.87	1.29	6.81	16.28	< 0.001
	Sex	124.90	1.60	29.16	2.97	0.006
	Context	61.76	1.51	186.68	-0.41	0.683

LMMs were performed on log-transformed response variables. Back-transformed estimates and standard errors (SE) are reported. The reference sex is “female” and the reference social context is “same-sex”. All models included mouse ID, observer ID, trial number, and observation period as random effects. Statistically significant results are shown in bold. Corresponds to Figure 3I.




Figure S1 | Comparison of individual and pair burrowing behavior in large, sand-filled enclosures. a) Number of burrows dug in individual (I) and pair (P) trials in three *Peromyscus* species. **b)** Maximum burrow length dug per trial by individuals and pairs of mice. Data are plotted on a square-root scale. Data points represent the mean of 1-7 (average 1.6) trials per individual or unique pair. * $P < 0.05$, *** $P < 0.001$. Error bars represent s.e.m.




Figure S2 | *P. maniculatus* females dig longer burrows when pregnant. **a**, Experimental design schematic. Individual females were first tested as virgins to determine baseline burrowing output (top). After a period of cohabitation with a conspecific male (middle), females were tested again (bottom). By chance, 19/36 females became pregnant during the cohabitation period. **b**, Maximum burrow length dug by females before (left) and after (right) male cohabitation. We found a significant interaction between cohabitation and pregnancy on the length of the longest burrow dug per trial (LMM, cohabitation: $F = 1.98$, $P = 0.162$, pregnancy: $F = 1.71$, $P = 0.200$, cohabitation*pregnancy: $F = 7.57$, $P = 0.007$). Using planned contrasts, we found an increase in burrowing output for pregnant females only (pregnant: $t = 3.01$, $P = 0.003$, non-pregnant: $t = 0.93$, $P = 0.355$). 15/19 (79%) pregnant females dug longer burrows after cohabitation with a male, with a median increase of 21% over their previous trials (virgin: 10.9 ± 0.9 cm, post-cohabitation: 13.6 ± 1.1 cm). By contrast, only 6/17 (35%) non-pregnant females dug longer burrows after cohabitation with a male. Each line represents the mean of 2 trials per individual, per timepoint. *** $P < 0.001$. Error bars represent s.e.m.




Figure S3 | Summary of *P. polionotus* burrowing behavior in narrow, transparent enclosures. **a)** Number of burrows dug in individual (I) and pair (P) trials. **b)** Maximum burrow length dug per trial by individuals and pairs of mice. **c)** Maximum burrow length dug per trial, broken down by trial-type. Data points in **a-c** represent the mean of 2 trials per individual or unique pair. **d)** Division of labor in same-sex (S, same) and opposite-sex (opp) pairs. **e)** Total digging duration for same-sex (FF, MM) and opposite-sex (FM) pairs, per ten-minute observation period. Data points in **d-e** represent the mean of 4 observations per pair, over 2 trials. * $P < 0.05$, *** $P < 0.001$. Error bars represent s.e.m.




Figure S4 | Sex differences in individual-level burrowing behavior in *P. polionotus*. **a-c**, Underground and digging behavior, pooled across same-sex and opposite-sex trials, for individual female ($n = 16$) and male ($n = 16$) mice. Behavior labels are as follows: “under” (i.e., mouse is underground, in the burrow), “extending” (i.e., mouse is digging at the leading edge of the burrow), “widening” (i.e., mouse is expanding the interior of the burrow), “hind-kick” (i.e., a powerful, expulsive kick that expels loose sand from the burrow), “obscured” (i.e., mouse is underground, but not visible). All behaviors (except hind-kicks) were scored as state events. Hind-kicking bouts were scored as point events. **a**, Number of bouts per observation period. **b**, Mean bout duration, calculated per observation period. **c**, Total duration of behavior per observation period. Data points represent the mean of 8 ten-minute observation periods per individual. Error bars represent s.e.m.